

BASE CHANGE, TRANSITIVITY AND KÜNNETH FORMULAS FOR THE QUILLEN DECOMPOSITION OF HOCHSCHILD HOMOLOGY

CHRISTIAN KASSEL and ARNE B. SLETSJØE

Let A be any commutative algebra over a commutative ring k and let M be any symmetric A -bimodule. In [Q], §8, Quillen proved that the Hochschild groups

$$H_*(A, M) = \text{Tor}_*^{A \otimes_k A}(M, A)$$

have a natural decomposition, called the Quillen decomposition,

$$H_n(A, M) \simeq \bigoplus_{p+q=n} D_q^{(p)}(A/k, M)$$

under the hypothesis that A is flat over k , containing the field \mathbb{Q} of rational numbers. The right-hand side is defined in terms of exterior powers of the cotangent complex of A over k . For $p = 1$, the groups $D_*^{(1)}(A/k, M)$ are isomorphic to the André-Quillen homology groups $D_*(A/k, M)$.

The purpose of this note is to prove base change, transitivity and Künneth formulas for all $D_*^{(p)}(A/k, M)$ – and hence for Hochschild homology in characteristic zero – extending analogous formulas established by André [A] and Quillen [Q] for $D_*(A/k, M)$.

Lately M. Ronco [R] proved that the Quillen decomposition coincides with a decomposition introduced by combinatorial methods on the level of Hochschild standard complex by Gerstenhaber-Schack [GS]. The latter decomposition coincides with another one due to Feigin-Tsygan [FT] and Burghlea-Vigué [BV] [V]. In the notation of [L], M. Ronco’s result can be written as follows (for all p and n)

$$D_{n-p}^{(p)}(A/k, M) \simeq H_n^{(p)}(A, M)$$

We assume all rings to be commutative with unit.

1. Definition of $D_*^{(p)}(A/k, M)$.

For any map of rings $u: k \rightarrow A$ and any nonnegative integer p , we define the simplicial A -module

$$\mathbf{L}_{A/k}^p = \Omega_{P/k}^p \otimes_P A$$

where P is a simplicial cofibrant k -algebra resolution of A in the sense of [Q]. By [Q], the simplicial A -module $\mathbf{L}_{A/k}^p$ is independent, up to homotopy equivalence, of the choice of P . In Quillen's notation

$$\mathbf{L}_{A/k}^p = \Lambda_A^p \mathbf{L}_{A/k}^1$$

where $\mathbf{L}_{A/k}^1$ is the cotangent complex. Thus we define

$$D_*^{(p)}(A/k, M) = H_* (\mathbf{L}_{A/k}^p \otimes_A M) \quad \text{and} \quad D_{(p)}^*(A/k, M) = H^*(\text{Hom}_A(\mathbf{L}_{A/k}^p, M))$$

for any A -module M .

REMARK 1.1. a) If $p = 0$, then $\mathbf{L}_{A/k}^p \simeq A$ and

$$D_n^{(0)}(A/k, M) = \begin{cases} M & \text{if } n = 0 \\ 0 & \text{otherwise.} \end{cases}$$

b) If $p = 1$, $D_*^{(1)}(A/k, M) = D_*(A/k, M)$ where the right-hand side was defined by André [A] and Quillen [Q]. These groups coincide with the Harrison groups [H] in characteristic zero.

We derive now some properties of the group $D_*^{(p)}(A/k, M)$ which are immediate consequences of Quillen's formalism.

LEMMA 1.2. $\mathbf{L}_{A/k}^p$ is a free simplicial A -module.

PROOF. This follows from the fact that if P is free over k , say $P = S_k(V)$, then

$$\Omega_{P/k}^p \otimes_P A \simeq (\Lambda_k(V) \otimes_k P) \otimes_P A \simeq \Lambda_k(V) \otimes_k A$$

COROLLARY 1.3. For any exact sequence of A -modules

$$0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$$

there are long exact sequences

$$\dots \rightarrow D_n^{(p)}(A/k, M') \rightarrow D_n^{(p)}(A/k, M) \rightarrow D_n^{(p)}(A/k, M'') \rightarrow D_{n-1}^{(p)}(A/k, M') \rightarrow \dots$$

and

$$\dots \rightarrow D_{(p)}^n(A/k, M') \rightarrow D_{(p)}^n(A/k, M) \rightarrow D_{(p)}^n(A/k, M'') \rightarrow D_{(p)}^{n+1}(A/k, M') \rightarrow \dots$$

The module $\mathbf{L}_{A/k}^p$ has the following vanishing property.

PROPOSITION 1.4. *If A is a free k -algebra, then $L_{A/k}^p$ has the homotopy type of $\Omega_{A/k}^p$. Consequently, for any A -modulus M*

$$D_n^{(p)}(A/k, M) = D_{(p)}^n(A/k, M) = 0 \quad \text{if } n \geq 1$$

and

$$D_0^{(p)}(A/k, M) = \Omega_{A/k}^p \otimes_A M \quad \text{and} \quad D_{(p)}^0(A/k, M) = \text{Hom}_A(\Omega_{A/k}^p, M)$$

PROOF. Take $P = A$.

2. Base change and Künneth formulas.

The following result states how L^p behaves under tensor products.

THEOREM 2.1. *If A and B are k -algebras such that $\text{Tor}_q^k(A, B) = 0$ for $q > 0$, then we have the following isomorphisms*

a) *Base change*

$$L_{A \otimes_k B/A}^p \simeq A \otimes_k L_{B/k}^p$$

b) *Künneth-type formula*

$$L_{A \otimes_k B/k}^p \simeq \bigoplus_{q+r=p} (L_A^q \otimes_k L_{B/k}^r)$$

PROOF. Under the hypothesis of theorem, if P (resp. Q) is a cofibrant k -resolution of A (resp. of B), then $A \otimes_k Q$ (resp. $P \otimes_k Q$) is a cofibrant resolution of $A \otimes_k B$ over A (resp. over k). Now

$$\begin{aligned} \Omega_{A \otimes_k Q/k}^p \otimes_{A \otimes_k Q} (A \otimes_k B) &\simeq (A \otimes_k \Omega_{Q/k}^p) \otimes_{A \otimes_k Q} (A \otimes_k B) \\ &\simeq A \otimes_k (\Omega_{Q/k}^p \otimes_Q B) \end{aligned}$$

For the Künneth formula, we have

$$\begin{aligned} \Omega_{P \otimes_k Q/k}^p \otimes_{P \otimes_k Q} (A \otimes_k B) &= \bigoplus_{q+r=p} ((\Omega_{P/k}^q \otimes_k \Omega_{Q/k}^r) \otimes_{P \otimes_k Q} (A \otimes_k B)) \\ &\simeq \bigoplus_{q+r=p} ((\Omega_{P/k}^q \otimes_P A) \otimes_k (\Omega_{Q/k}^r \otimes_Q B)) \end{aligned}$$

COROLLARY 2.2. *Under the same hypotheses as Theorem 2.1, and for any $A \otimes_k B$ -module M , we have the following isomorphisms of graded modules*

$$D^{(p)*}(A \otimes_k B/A, M) \simeq D_*^{(p)}(B/k, M)$$

and

$$D_*^{(p)}(A \otimes_k B/k, M) \simeq \bigoplus_{q+r=p} D_*^{(q)}(A/k, M) \otimes_k D_*^{(r)}(B/k, M)$$

In characteristic zero the corresponding isomorphism for $\mathrm{HH}_*^{(p)}(A \otimes_k B)$ and for the cyclic groups $\mathrm{HC}_*^{(p)}(A \otimes_k B)$ are also proved in [K].

3. Transitivity.

Suppose we have maps $k \xrightarrow{u} A \xrightarrow{v} B$ of commutative rings. We start by defining a filtration of $\Omega_{B/k}^p$. Let $F_A^i(\Omega_{B/k}^p)$ be the sub- A -module of $\Omega_{B/k}^p$ generated by $b_0 db_1 \dots db_p$ where at least i elements among b_1, \dots, b_p lie in A . We have the following sequence of inclusions of A -modules,

$$\Omega_{B/k}^p = F_A^0 \supset F_A^1 \supset \dots \supset F_A^p = \Omega_{A/k}^p \otimes_k B$$

LEMMA 3.1. *If B is A -free and A is k -free, then the map*

$$\psi_i: \Omega_{A/k}^i \otimes_A \Omega_{B/A}^{p-i} \rightarrow F_A^i / F_A^{i+1}$$

given by

$$\psi(a_0 da_1 \dots da_i \otimes b_0 db_{i+1} \dots db_p) = a_0 b_0 da_1 \dots da_i \cdot db_{i+1} \dots db_p$$

is an isomorphism.

PROOF. First check that ψ_i is well-defined without any hypothesis on A and B . If $A = S_k(V)$ and $B = S_A(A \otimes W) = S_k(V) \otimes S_k(W) = S_k(V \oplus W)$ one computes easily both source and target of ψ_i .

THEOREM 3.2. *Let $k \xrightarrow{u} A \xrightarrow{v} B$ be maps of commutative rings and let M be a B -module. Then there is a spectral sequence (E^r, d^r) converging to $D_*^{(p)}(B/k, M)$. The k -module $E_{i,j}^1$ have the following properties:*

- a) $E_{i,j}^1 = 0$ for $i > 0$ or $i < -p$.
- b) $E_{0,j}^1 = D_j^{(p)}(B/A, M)$ and $E_{-p,j}^1 = D_{j-p}^{(p)}(A/k, M)$.
- c) Fix any p . For every i there is a first quadrant spectral sequence $({}^{(i)}E^r, d^r)$ converging to $E_{-i,i+*}^1$ such that

$${}^{(i)}E_{k,l}^2 = D_k^{(i)}(A/k, D_l^{(p-i)}(B/A, M))$$

REMARK 3.3. a) The edge homomorphisms

$$D_j^{(p)}(B/k, M) \rightarrow E_{0,j}^1 = D_j^{(p)}(B/A, M)$$

and


$$E_{-p,p+j}^1 = D^{(p)}(A/k, M) \rightarrow D_j^{(p)}(B/k, M)$$

are the natural homomorphisms. For $p = 1$, the first spectral sequence reduces to two columns, so that one recovers the well-known long exact sequence

$$\dots \rightarrow D_j(A/k, M) \rightarrow D_j(B/k, M) \rightarrow D_j(B/A, M) \rightarrow D_{j-1}(A/k, M) \rightarrow \dots$$

b) Applying Theorem 3.2 to the map of rings $k \rightarrow A \rightarrow A \otimes_k B$, one sees that the spectral sequences degenerate and one recovers the Künneth formula of Corollary 2.2.

PROOF OF THEOREM 3.2. Let P be a simplicial cofibrant k -resolution of A . Consider the composite map $P \rightarrow A \rightarrow B$ and choose a simplicial cofibrant P -resolution Q of B . Let us consider the following commutative diagram



Then it follows from [Q] that $A \otimes_P Q$ is a simplicial cofibrant A -resolution of B .

We apply the construction of Lemma 3.1 to the map of rings $k \rightarrow P \rightarrow Q$. Then we get a filtration of $\Omega_{Q/k}^p \otimes_Q M$ such that the associated graded is $\Omega_{P/k}^i \otimes_P \Omega_{Q/P}^{p-i} \otimes_Q M$. This yields the first spectral sequence with

$$E_{i,j}^1 = H_{i+j}(\Omega_{P/k}^i \otimes_P (\Omega_{Q/P}^{p-i} \otimes_Q M))$$

converging to $H_{i+j}(\Omega_{Q/k}^p \otimes_Q M)$ which is $D_{i+j}^{(p)}(B/k, M)$ because Q is also a simplicial cofibrant k -resolution of B .

To compute the homology of $\Omega_{P/k}^i \otimes_P \Omega_{Q/P}^{p-i} \otimes_Q M$ we use the fact that it has a double simplicial structure. Therefore it gives rise to a spectral sequence with E^2 -term of the form

$$\begin{aligned} {}^{(i)}E_{k,1}^2 &= H_k(\Omega_{P/k} \otimes_P H_1(\Omega_{Q/P}^{p-i} \otimes_Q M)) \\ &= D_k^{(i)}(A/k, H_1(\Omega_{Q/P}^{p-i} \otimes_Q M)) \end{aligned}$$

Now we use the base change formula of Theorem 2.1 to get the following isomorphism of P -modules

$$\begin{aligned} D_i^{(p-i)}(B/A, M) &= H_i(\Omega_{A \otimes_P Q/A}^{(p-i)} \otimes_{A \otimes_P Q} M) \\ &= H_i(\Omega_{Q/P}^{(p-i)} \otimes_Q M) \end{aligned}$$

4. Applications.

The following is an extension of Quillen’s Theorem 5.4 [Q].

PROPOSITION 4.1. Assume that $k \supset \mathbb{Q}$ and $\Omega_{A/k}^1$ is A -flat.

- i) If $\text{Spec } A \rightarrow \text{Spec } k$ is étale, then $L_{A/k}^p$ is acyclic for $p \geq 1$.
- ii) If $\text{Spec } A \rightarrow \text{Spec } k$ is smooth, then $L_{A/k}^p \simeq \Omega_{A/k}^p$.

PROOF. i) Let P be a simplicial cofibrant k -resolution of A . By [Q], if A is étale over k , then $\Omega_{P/k}^1 \otimes_P A = \mathbf{L}_{A/k}^1$ is acyclic. Hence

$$\mathbf{L}_{A/k}^p = \mathbf{L}_A^p \mathbf{L}_{A/k}^1$$

which is a direct summand (in characteristic zero) of $(\mathbf{L}_{A/k}^1)^{\otimes p}$ is acyclic.

ii) We have the following isomorphisms

$$\mathbf{L}_{A/k}^p = \mathbf{L}_P^p \Omega_{P/k}^1 \otimes_P A \simeq \mathbf{L}_A^p \Omega_{A/k}^1 \otimes_A A \simeq \Omega_{A/k}^p$$

in the derived category of A -modules.

COROLLARY 4.2. *Under the hypothesis of Proposition 4.1 and if A is smooth over k , then for all p*

$$D_n^{(p)}(A/k, M) = \begin{cases} \Omega_{A/k}^p \otimes_A M & \text{if } n = 0 \\ 0 & \text{otherwise} \end{cases}$$

SPECIAL CASES 4.3. Let $k \rightarrow A \rightarrow B$ be maps of rings such that $k \supset \mathbf{Q}$ and let M be a B -module.

a) If A is smooth over k , then by Theorem 3.2 and Corollary 4.2 the spectral sequence converging to $D_*^{(p)}(B/k, M)$ has E^1 -term given by

$$E_{-i, i+j}^1 = \Omega_{A/k}^i \otimes_A D_j^{(p-i)}(B/A, M)$$

b) If A/k is étale, we get: $D_*^{(p)}(B/k, M) = D_*^{(p)}(B/A, M)$ from Theorem 3.2 and Prop. 4.1. i. The resulting isomorphism for Hochschild homology

$$H_*(B/k, M) \simeq H_*(B/A; M)$$

was proved by Gerstenhaber-Schack [GES].

c) If B is smooth over A , then the E^1 -terms are given by

$$E_{-i, i+j}^1 = D_j^{(i)}(A/k, \Omega_{B/A}^{(p-i)} \otimes_B M)$$

If moreover B is étale over A , then $\Omega_{B/A}^p = 0$ for $p > 0$. From Theorem 3.2 we get the following isomorphism:

$$D_*^{(p)}(B/k, M) \simeq D_*^{(p)}(A/k, M)$$

If the B -module M is extended from A , i.e. is of the form $B \otimes_A N$ where N is an A -module, then we have the following étale descent isomorphism

$$D_*^{(p)}(B/k, M) \simeq D_*^{(p)}(A/k, N) \otimes_A B$$

When $N = A$, we thus recover Theorem 0.1 of [WG] stating that

$$H_*(B, B) \simeq H_*(A, A) \otimes_A B$$

REFERENCES

- [A] M. André, *Homologie des algèbres commutatives*, Springer Gru. 206, Berlin-Heidelberg-New York, 1974.
- [BV] D. Burghelea and M. Vigué-Poirrier, *Cyclic homology of commutative algebras I*, Lect. Notes in Math. 1318 (1988), 51–72.
- [FT] B. L. Feigin and B. L. Tsygan, *Additive K-theory and crystalline cohomology*, Functional Anal. Appl. 19 (1985), 124–132.
- [GES] M. Gerstenhaber and S. D. Schack, *Relative Hochschild cohomology, rigid algebras and the Bockstein*, J. Pure Appl. Algebra 43 (1986), 53–74.
- [GS] M. Gerstenhaber and S. D. Schack, *A Hodge-type decomposition for commutative algebra cohomology*, J. Pure Appl. Algebra 48 (1987), 229–247.
- [H] D. K. Harrison, *Commutative algebras and cohomology*, Trans. Amer. Math. Soc. 104 (1962), 191–204.
- [K] C. Kassel, *Une formule de Künneth pour la décomposition de l'homologie cyclique des algèbres-commutatives*, Preprint, Strasbourg (1989).
- [L] J.-L. Loday, *Opérations sur l'homologie cyclique des algèbres commutatives*, Invent. Math. 96 (1989), 205–230.
- [Q] D. Quillen, *On the (co)homology of commutative rings*, Proc. Sympos. Pure Math. 17 (1970), 65–87.
- [R] M. Ronco, *Sur l'homologie d'André-Quillen*, Preprint IRMA, Strasbourg (1990).
- [V] M. Vigué-Poirrier, *Décomposition de l'homologie cyclique des algèbres différentielles graduées commutatives*, to appear in K-Theory.
- [WG] C. A. Weibel and S. C. Geller, *Étale descent for Hochschild and cyclic homology*, Preprint (1990), to appear in Comment Math. Helv.

UNIVERSITÉ LOUIS PASTEUR
DÉPT. DE MATHÉMATIQUE
7, RUE RENÉ DESCARTES
F-67084 STRASBOURG CEDEX
FRANCE

MATEMATISK INSTITUTT
UNIVERSITETET I OSLO
PB. 1053 BLINDERN
N-0316 OSLO 3
NORWAY