

# BOUNDED LINEAR FUNCTIONALS ON VECTOR-VALUED $H^\infty$ SPACES

WOLFGANG HENSGEN

## Introduction.

The theory of the space  $H^\infty$ , its dual, and its predual  $L^1/H_0^1$ , is known to be difficult (see e.g. [Bo]). In this work I transfer some of the easier parts of this theory to a vector resp. operator-valued context.

Let  $X$  be a complex Banach space and  $H^\infty(X)$  [ $H^\infty(X)$ ] the space of bounded holomorphic  $X$ -valued functions on the unit disc [admitting strong radial boundary values a.e.]. By the simple but for this paper crucial lemma 2.1 I can prove the F. and M. Riesz theorem for  $H^\infty(X)$  (2.2). Immediate consequences are that for reflexive  $X$  the spaces  $H^1(X)$  and  $L^1(X)/H_0^1(X)$  are  $L$ -embedded and  $C(X)/A(X)$  is an  $M$ -ideal in the bidual (2.4–2.6). In §3, I study integral resp. strongly integral functionals on  $H^\infty(X)$ , i.e. those given by Gel'fand integrable functions  $g \in L^1(X', X)$  resp. Bochner integrable functions  $g \in L^1(X')$ . I found it interesting to see how the various characterizations of integral functionals in the scalar case [Kh 2; Ga] group exactly into two sets of conditions in the vector case: the first set, formulated on the circle and describing the integral functionals (3.1), and the second group, formulated on the disc and singling out the strongly integral functionals (3.5). Contrary to the scalar situation, some locally convex theory (Grothendieck's dual characterization of completeness) enters the proof of the latter theorem. The Banach space theoretic consequence of the former is that  $L^1(X)/H_0^1(X)$  has property (X) if  $X$  is reflexive (3.2).

The paper concludes in §4 with spaces of operator-valued functions. Let  $K, N, B$  denote the spaces of compact resp. nuclear resp. bounded operators on separable Hilbert space. I prove an F. and M. Riesz theorem for  $H^\infty(B)$  (4.5); consequently  $H^1(N)$  and  $L^1(N)/H_0^1(N)$  are  $L$ -embedded and  $C(K)/A(K)$  is an  $M$ -ideal in the bidual (4.6, 4.7). The result on  $L^1(N)/H_0^1(N)$  is related to a question of Pisier. Moreover,  $L^1(N)/H_0^1(N)$  enjoys property (X) (4.8).

§§1–3 of the paper are essentially a part of the author's habilitation thesis [He 3]. I would like to thank Professor A. V. Bukhvalov for his suggestions and stimulations to do this work. §4 results from a couple of stimulating discussions with Dr. H. Pfitzner who in particular pointed out Pisier's question to me. 4.8 answers a question of the referee.

## §1. Preliminaries.

The basic theory of vector-valued Hardy spaces as developed mainly by Bukhvalov and Danilevich [Bu 1; BD] is assumed familiar. See [He 1; 3] for detailed and [LM] for a very concise presentation. I fix some notation.

1.1. The basic measure space is the circle group  $\mathbb{T}$  with normalized Haar measure  $\lambda$ . For  $1 \leq p \leq \infty$ ,  $L^p(\lambda; X) = L^p(X)$  is the  $X$ -valued Lebesgue-Bochner space and  $L^p(\lambda; X', X) = L^p(X', X) (= L^p(X')$  iff  $X'$  has RNP) denotes the space of  $X$ -scalar equivalence classes of weak\* measurable functions  $f: \mathbb{T} \rightarrow X'$  with  $|f| := L^0 - \sup_{x \in B_x} |\langle x, f \rangle| \in L^p$ , normed by  $\|f\|_p = \|\ |f|\ \|_p$ . Such a function is Gel'fand integrable [DU, p. 53]. Under the canonical dual pairing,  $L^\infty(X', X)$  is the dual of  $L^1(X)$ .

1.2. The dual of  $L^\infty(X)$  has been described by various authors, see [CV, VII; BL]. Any member  $g \in L^1(X', X)$  defines a functional  $\varphi_g \in L^\infty(X)'$  by the formula  $\varphi_g(f) := \int \langle f, g \rangle d\lambda$ ; moreover  $\|\varphi_g\| = \|g\|_1$ . (Proof for " $\geq$ ":  $\|g\|_1 =$  total variation of the vector measure  $g \cdot \lambda$  [He 1, (0.8)2°]  $\leq \|\varphi_g\|$ , by definition of the variation.) The functionals  $\varphi_g$  are called *integral* and the space of integral functionals is identified with  $L^1(X', X)$ .

A functional  $\varphi \in L^\infty(X)'$  is called *concentrated* on a (measurable) set  $E \subset \mathbb{T}$  if  $\varphi(f) = \varphi(\mathbf{1}_E f) \forall f \in L^\infty(X)$ , and *singular* if concentrated on sets of arbitrarily small  $\lambda$ -measure. The space of singular functionals is denoted by  $L^\infty(X)'_s$ .

*Vector-valued Yosida-Hewitt decomposition:*

$$L^\infty(X)' = L^1(X', X) \oplus_1 L^\infty(X)'_s, (\oplus_1: L\text{-decomposition.})$$

1.3. The vector-valued Hardy spaces used in this paper are:  $H^p(X) := \{f \in L^p(X): \hat{f}(n) = 0 \forall n < 0\} \cong \{f: D \rightarrow X \text{ holomorphic: } \|f\|_p := \sup_{r < 1} \|f_r\|_p < \infty \text{ and } \lim_{r \rightarrow 1} f(re^{i\theta}) \text{ exists a.e.}\}$ ;  $H^p(X') := \{f \in L^p(X', X): \hat{f}(n) = 0 \forall n < 0\} \cong \{f: D \rightarrow X' \text{ holomorphic: } \|f\|_p := \sup_{r < 1} \|f_r\|_p < \infty\}$  ( $= H^p(X')$  iff  $X'$  has ARNP);  $A(X) := \{f \in C(\mathbb{T}; X): \hat{f}(n) = 0 \forall n < 0\}$  ( $X$ -valued disc "algebra"). With the usual meaning of the subscript 0, the following canonical dualities hold:

$$(C(X)/A(X))' = H_0^1(X'); H_0^1(X)' = L^\infty(X', X)/H^\infty(X'); (L^1(X)/H_0^1(X))' = H^\infty(X').$$

## §2. F. and M. Riesz Theorem for $H^\infty(X)$ .

The following lemma exploits the  $L^\infty$  module structure of  $L^\infty(X)$  by “contracting” a functional with a function.

2.1. LEMMA. *Let  $h \in L^\infty(X)$  be fixed. For  $\varphi \in L^\infty(X)'$  define  $\varphi h \in L^{\infty'}$  by the formula  $(\varphi h)(u) := \varphi(uh)$ ,  $u \in L^\infty$ . If  $\varphi = \varphi_i + \varphi_s$  is the vector-valued Yosida-Hewitt decomposition (1.2) of  $\varphi \in L^\infty(X)'$  then  $\varphi h = \varphi_i h + \varphi_s h$  is the (scalar) Yosida-Hewitt decomposition of  $\varphi h \in L^{\infty'}$ . (In other words,  $(\varphi h)_i = \varphi_i h$  and  $(\varphi h)_s = \varphi_s h$ .)*

PROOF. Let  $g \in L^1(X', X)$  be the function defining the integral part  $\varphi_i$ , i.e.  $\varphi_i(f) = \int \langle f, g \rangle d\lambda \quad \forall f \in L^\infty(X)$ . Then  $(\varphi_i h)(u) = \int \langle uh, g \rangle d\lambda = \int u \langle h, g \rangle d\lambda \quad \forall u \in L^\infty$ , so that  $\varphi_i h$  is integral, given by  $\langle h, g \rangle \in L^1$ . On the other hand, for an arbitrary  $\Psi \in L^\infty(X)'$ , the contraction  $\Psi h \in L^{\infty'}$  is concentrated on every set on which  $\Psi$  is concentrated. In particular,  $\varphi_s h$  is singular together with  $\varphi_s$ .

2.2. F. AND M. RIESZ THEOREM FOR  $H^\infty(X)$ : *Let  $\varphi = \varphi_i + \varphi_s$  be the vector-valued Yosida-Hewitt decomposition (1.2) of  $\varphi \in L^\infty(X)'$ . If  $\varphi|H_0^\infty(X) = 0$  then  $\varphi_i|H_0^\infty(X) = 0$  and  $\varphi_s|H^\infty(X) = 0$ . In particular, the annihilator  $H^\infty(X)^\perp$  is invariant under the Yosida-Hewitt projection  $P: L^\infty(X)' \rightarrow L^1(X', X)$ ,  $\varphi \mapsto \varphi_i$ .*

PROOF. In the scalar case, Gel'fand transformation translates the theorem into the well-known F. and M. Riesz theorem in logmodular algebras [Ho, p. 186; Ga, V. 4.4]. Turning to the vector case, let  $\varphi \in L^\infty(X)'$  be given with  $\varphi|H_0^\infty(X) = 0$ , and fix  $h \in H^\infty(X)$ ; one has to show  $\varphi_s(h) = 0$ . To this end, consider  $\varphi h \in L^{\infty'}$  as in lemma 2.1 and note that  $\varphi h|H_0^\infty = 0$ , exploiting the fact that  $u \in H_0^\infty$ ,  $h \in H^\infty(X)$  implies  $uh \in H_0^\infty(X)$ . By the scalar case,  $(\varphi h)_s|H^\infty = 0$ , whence by the lemma,  $\varphi_s(h) = \varphi_s h(1) = (\varphi h)_s(1) = 0$ .

If one is interested only in the invariance result then, for reflexive  $X$ , a completely different approach is possible, discovered by Godefroy [Go 1] (for the scalar case). It is based on (a vector-valued version of) the Bukhvalov-Lozanovskii theorem [Bu 2; BL] and thus lastly on vector lattices instead of uniform algebras. Still another (scalar) proof, via the notion of “strictly convergent” (=  $wuC$ ) series in  $L^\infty$ , has been developed by Barbey and König [BK, VIII.]. See also Ando [A, 4.].

2.3. COROLLARY. *If  $\varphi \in L^\infty(X)'$  is singular and  $\varphi|H^\infty(X) = 0$  then  $\varphi|C(X) = 0$ .*

PROOF. Apply the usual iteration [Ho, p. 186 f.]: The functional  $f \mapsto \varphi(e^{-i\theta} f)$  on  $L^\infty(X)$  vanishes on  $H_0^\infty(X)$  and is also singular, hence by 2.2:  $\varphi(e^{-i\theta} x) = 0$  for all  $x \in X$ , etc. Thus  $\varphi$  vanishes on all trigonometric polynomials which are dense in  $C(X)$ .

2.4. PROPOSITION. *If  $X$  is reflexive then  $H_0^1(X)'' = H_0^1(X) \oplus_1 (L^\infty(X')'_s \cap H^\infty(X')^\perp)$ . Thus  $H_0^1(X)$  (hence also  $H^1(X)$ ) is  $L$ -embedded in its bidual.*

PROOF. Using reflexivity all the time, 1.3 and 1.2 yield  $H_0^1(X)'' = (L^\infty(X')/H^\infty(X'))' = H^\infty(X')^\perp \subset L^\infty(X') = L^1(X) \oplus_1 L^\infty(X')'_s$ . Trivially, the restriction of an  $L$ -projection  $P$  to an invariant subspace  $Y$  is an  $L$ -projection in  $Y$  onto  $Y \cap \text{range}(P)$ . With  $Y := H^\infty(X')^\perp$ , 2.2 implies  $H_0^1(X)'' = (L^1(X) \cap H^\infty(X')^\perp) \oplus_1 (L^\infty(X')'_s \cap H^\infty(X')^\perp)$ , and the first summand equals  $H_0^1(X)$ .

2.5. PROPOSITION. *If  $X$  is reflexive then  $C(X)/A(X)$  is an  $M$ -ideal in its bidual  $L^\infty(X)/H^\infty(X)$ . (See [HWW] for definition and consequences – e.g., property (u).)*

PROOF. It is convenient to prove the assertion for  $X'$ , so let  $Z := C(X')/A(X')$ ; one has to show the  $L$ -nature of the canonical decomposition  $Z''' = Z' \oplus Z^\perp$ . Now by 1.3,  $Z' = H_0^1(X)$  and this decomposition reads  $H_0^1(X)'' = H_0^1(X) \oplus Z^\perp$ . On the other hand, 2.4 says  $H_0^1(X)'' = H_0^1(X) \oplus_1 (L^\infty(X')'_s \cap H^\infty(X')^\perp)$ . I claim that this last  $L$ -summand is contained in  $Z^\perp$  (then it will coincide with  $Z^\perp$  by pure linear algebra): The annihilator  $Z^\perp$  of  $Z := C(X')/A(X')$  in  $Z''' = H_0^1(X)'' = H^\infty(X')^\perp \subset L^\infty(X')$  equals  $\{\varphi \in H^\infty(X')^\perp : \varphi|C(X') = 0\}$  and the claim follows from 2.3.

This proof differs from Luecking's (scalar) proof [HWW, p. 109] in the use of 2.4. If this is avoided as in [HWW, end of p. 109], one gets 2.4 as a consequence.

2.6. PROPOSITION. *If  $X$  is reflexive then  $L^1(X)/H_0^1(X)$  is  $L$ -embedded in its bidual.*

PROOF. By 1.3, 1.2 and reflexivity,  $(L^1(X)/H_0^1(X))'' = H^\infty(X')' = L^\infty(X')/H^\infty(X')^\perp = (L^1(X) \oplus_1 L^\infty(X')'_s)/H^\infty(X')^\perp$ . It is well-known and not hard to prove (see e.g. [He 3, 2.3], [HWW, I.1.15]) that an  $L$ -projection  $P$  in a space  $Z$  canonically induces an  $L$ -projection  $\bar{P}$  in  $Z/Y$  for a  $P$ -invariant subspace  $Y$ , with  $\text{range}(\bar{P}) = \text{range}(P)/(Y \cap \text{range}(P))$  (as canonically isometrically embedded in  $Z/Y$ ). With  $Y := H^\infty(X')^\perp$ , 2.2 implies  $(L^1(X)/H_0^1(X))'' = L^1(X)/(L^1(X) \cap H^\infty(X')^\perp) \oplus_1 \dots = L^1(X)/H_0^1(X) \oplus_1 \dots$

It follows from 2.6 and Godefroy's result [Go 1, 4; HWW, IV.2.2] that  $L^1(X)/H_0^1(X)$  is weakly sequentially complete for reflexive  $X$ . This fact (and probably 2.6) was obtained first by Petrenko [Bu 2, 5.3], cf. [He 2]. Recently, Pfitzner [Pf 1; HWW, IV.2.7] proved that spaces  $L$ -embedded in the bidual even enjoy Pelczyński's property ( $V^*$ ). For the case of  $L^1(X)/H_0^1(X)$  with  $X$  reflexive, property ( $V^*$ ) has been obtained earlier by this author [He 3, 3.6; He 4, 2.2] adapting Pelczyński's scalar proof [Pe, 7.1]. See also 3.2.

### §3. Integral and strongly integral functionals.

For the scalar-valued origins of the following theorem, see [Kh 2, Theorem 1; BK, VIII.3.1; Ga, V.5.3; Kh 1, p. 300].

3.1. THEOREM (Characterization of integral functionals). *For  $\varphi \in H^\infty(X)'$ , the following conditions are equivalent:*

1°  $\varphi$  is integral, i.e.  $\exists g \in L^1(X', X)$  with  $\varphi(f) = \int \langle f, g \rangle d\lambda \forall f \in H^\infty(X)$

2°  $f_n \in B_{H^\infty(X)}$  and  $f_n(e^{it}) \rightarrow 0$  a.e. on  $T$  implies  $\varphi(f_n) \rightarrow 0$

3°  $f_n \in H^\infty(X)$ ,  $\sum_{n=1}^{\infty} \|f_n(e^{it})\| \leq 1$  a.e. on  $T$  (whence  $\sum_{n=1}^{\infty} f_n(e^{it})$  is a member of  $H^\infty(X)$ )

implies  $\varphi\left(\sum_{n=1}^{\infty} f_n\right) = \sum_{n=1}^{\infty} \varphi(f_n)$

4°  $\varphi|_{B_{H^\infty(X)}}$  has a point of continuity for  $\|\cdot\|_1$ .

PROOF. 1°  $\Rightarrow$  2°:  $|\langle f_n, g \rangle| \leq \|f_n(\cdot)\| \|g\| \rightarrow 0$  a.e. on  $T$  (see 1.1 for notation and [He 1, (0.5) 5°] for the estimate), dominated by  $|g| \in L^1$ . Apply Lebesgue's theorem.

2°  $\Rightarrow$  3°: As regards the assertion in parentheses, clearly  $\sum_{n=1}^{\infty} f_n(e^{it})$  converges in  $X$  a.e. on  $T$ ,  $\sum_{n=1}^{\infty} f_n \in L^\infty(X)$  and  $\left(\sum_{n=1}^{\infty} f_n\right)^\wedge(k) = 0 \forall k < 0$  by bounded convergence.

Now apply 1° to the remainders  $\sum_{k=n}^{\infty} f_k(e^{it})$ .

3°  $\Rightarrow$  1°: In the scalar case, various proofs have been developed: [Ga, V.5.3; Go 1, 33; Go 2, V.4(5); BK, VIII.3.1]. The vector case is handled by lemma 2.1. Let  $\varphi \in L^\infty(X)'$  satisfy 3°. Since the integral part of  $\varphi$  (1.2) satisfies 3°, one can assume  $\varphi$  singular and has to prove  $\varphi = 0$  on  $H^\infty(X)$ . So let  $h \in H^\infty(X)$  be given; the contraction  $\varphi h \in L^\infty'$ :  $u \mapsto \varphi(uh)$  is singular by 2.1 and respects the convergence of

type 3° ( $X = \mathbb{C}$ ): if  $u_n \in H^\infty$ ,  $\sum_{n=1}^{\infty} |u_n(e^{it})| \leq 1$  a.e. then  $u_n h \in H^\infty(X)$ ,

$\sum_{n=1}^{\infty} \|u_n(e^{it})h(e^{it})\| \leq \|h\|_\infty$  a.e., whence by assumption  $\varphi h\left(\sum_{n=1}^{\infty} u_n h\right) = \varphi\left(\sum_{n=1}^{\infty} u_n h\right) =$

$\sum_{n=1}^{\infty} \varphi(u_n h) = \sum_{n=1}^{\infty} \varphi h(u_n)$ . By the scalar case  $\varphi h = 0$  on  $H^\infty$  and  $\varphi(h) = \varphi h(1) = 0$ .

2°  $\Rightarrow$  4°: It is very easily seen that actually  $2^\circ \Leftrightarrow \varphi|_{B_{H^\infty(X)}}$  is continuous for  $\|\cdot\|_1$ .

4°  $\Rightarrow$  1°: This rests on the Khavin lemma, with the aid of which Khavin [Kh 1, p. 299 f.] proved that if  $\varphi \in L^\infty'$  and  $\varphi|_{B_{H^\infty}}$  has a point of continuity for  $\|\cdot\|_1$  then the singular part of  $\varphi$  (1.2) vanishes on  $H^\infty$ . For the vector-valued case, the same proof works, and is given in detail in [He 2, 3.2].

NOTE. As in the scalar case,  $2^\circ \Rightarrow 1^\circ$  has an easier proof, transferring [Kh 2, p. 88] or [Ga, V Exercise 19] to the vector situation.

3.2. COROLLARY. *If  $X$  is reflexive then  $L^1(X)/H_0^1(X)$  has property (X) of Godefroy and Talagrand. Consequently,  $L^1(X)/H_0^1(X)$  is strongly unique predual of  $H^\infty(X')$ .*

PROOF. By definition, a Banach space  $Z$  has property (X) if  $z'' \in Z''$ ,  $z'' \left( \sum_{n=1}^\infty z'_n \right) = \sum_{n=1}^\infty z''(z'_n) \forall$  wuC series  $\sum z'_n$  in  $Z'$  implies  $z'' \in Z$  ( $\sum_{n=1}^\infty z'_n$  is taken for  $\sigma(Z', Z)$ ). Let  $\varphi \in (L^1(X)/H_0^1(X))'' = H^\infty(X')$  be given, and suppose  $\varphi \left( \sum_{n=1}^\infty f_n \right) = \sum_{n=1}^\infty \varphi(f_n) \forall$  wuC series  $\sum f_n$  in  $H^\infty(X')$ , summed for  $\sigma(H^\infty(X'), L^1(X)/H_0^1(X))$ . The "test series" of 3.1.3° ( $X'$ ) are wuC (since wuC =  $w^*u$ C in dual spces), and of course the pointwise limit considered there coincides with the weak\* limit. Thus 3.1 yields that  $\varphi$  is given by integration against a function  $g \in L^1(X)$ , i.e.  $\varphi \in L^1(X)/H_0^1(X)$ . The consequence is well-known [GT, 5; Go 2, V.3].

NOTES. i) (X) entails (V\*) [Go 2, V.5; GS, III.2; E, 13] providing still another proof of (V\*) for  $L^1(X)/H_0^1(X)$  if  $X$  is reflexive.

ii) In the scalar case, the unique predual result is due to Ando [A] and Wojtaszczyk [W], and property (X) of  $L^1/H_0^1$  goes back to Barbey [BK, VIII.].

3.3. REMARK. Let  $\mu$  be a finite measure and  $\varphi \in L^\infty(\mu; X')$ . Then the criterion analogous to 3.1.3° holds for  $\varphi$  to be integral. Namely, for  $X = \mathbb{C}$  this is the well-known property (X) of  $L^1(\mu)$  [Go 1, 30]; the general case is reduced to the scalar case by lemma 2.1 as before. Now the same proof as in 3.2 shows that for  $X$  reflexive,  $L^1(\mu; X)$  has property (X) and is hence strongly unique predual of  $L^\infty(\mu; X')$ . The latter conclusion follows also from [CG 2, Theorem 1] in connection with [CG 1, Remark 1].

3.4. DEFINITION. a) [Kh2, 5.] A continuous function  $w: [0, 1] \rightarrow \mathbb{R}_+$  with  $w(r) > 0$  if  $r < 1$ ,  $w(1) = 0$  is called a *weight*. For  $f: D \rightarrow X$ , put  $\|f\|_w := \sup_{x \in D} \|f(z)\| w(z)$  where  $w(z) := w(|z|)$ .

b) The locally convex topology on  $H^\infty(X)$  induced by the family of norms  $\|\cdot\|_w$  is the *strict topology*  $\beta$ .

It is easy to see that this is the vector-valued analogue of the strict topology on  $H^\infty$  studied in [RR; RS; Co, V.; BS].

3.5. THEOREM (Characterization of strongly integral functionals). *For  $\varphi \in H^\infty(X)$ , the following conditions are equivalent:*

$$1^\circ \varphi \text{ is strongly integral, i.e. } \exists g \in L^1(\mathbb{T}, \lambda; X') \text{ with } \varphi(f) = \int_{\mathbb{T}} \langle f, g \rangle d\lambda \forall f \in H^\infty(X)$$

$$2^\circ \exists g \in L^1(D, \lambda_2; X') \text{ with } \varphi(f) = \iint_D \langle f, g \rangle d\lambda_2 \quad \forall f \in H^\infty(X) \text{ } (\lambda_2: \text{area measure})$$

3°  $\exists$  weight  $w \exists C \in \mathbb{R}_+$  with  $|\varphi(f)| \leq C \|f\|_w \quad \forall f \in H^\infty(X)$  (i.o.w.,  $\varphi$  is  $\beta$ -continuous)

$$4^\circ \exists m \in M(D; X') \text{ with } \varphi(f) = \iint_D \langle f, dm \rangle \quad \forall f \in H^\infty(X) \text{ } (M: \text{vector measures}$$

$[\sigma$ -additive] of bounded variation)

$$5^\circ f_n \in B_{H^\infty(X)} \text{ and } f_n(z) \rightarrow 0 \quad \forall z \in D \text{ implies } \varphi(f_n) \rightarrow 0.$$

PROOF. First I show the equivalence of 1° through 4°, following [Kh 2, 5.] (cf. also [RS, 4.]). The proof rests on the following lemma on balayage whose proof is omitted (see [He 3, Lemma 2.8]).

$$\text{LEMMA. For } m \in M(D; X) \text{ put } B[m]: \mathbb{T} \rightarrow X, B[m](e^{i\theta}) := \iint_D P_z(\vartheta) m(dz) \text{ } (P_z(\vartheta):$$

Poisson kernel).

i)  $B[m](e^{i\theta})$  is defined a.e. on  $\mathbb{T}$  and  $B: M(D; X) \rightarrow L^1(\mathbb{T}; X)$  is an operator of norm  $\leq 1$

$$\text{ii) For } m \in M(D; X) \text{ and } f \in H^\infty(X'): \int_{\mathbb{T}} \langle B[m], f \rangle d\lambda = \iint_D \langle dm, f \rangle$$

iii)  $B: L^1(D, \lambda_2; X) \rightarrow L^1(\mathbb{T}, \lambda; X)$  is surjective.

1°  $\Rightarrow$  2°: For  $g \in L^1(\mathbb{T}; X')$  representing  $\varphi$  as in 1°, the lemma (iii) yields some  $\tilde{g} \in L^1(D; X')$  with  $B[\tilde{g}] = g$ . And by (ii), for  $f \in H^\infty(X) \subset H^\infty(X'')$ :

$$\varphi(f) = \int_{\mathbb{T}} \langle f, g \rangle d\lambda = \int_{\mathbb{T}} \langle f, B[\tilde{g}] \rangle d\lambda = \iint_D \langle f, \tilde{g} \rangle d\lambda_2.$$

$$2^\circ \Rightarrow 3^\circ: \text{ For } g \in L^1(D; X') \text{ representing } \varphi \text{ as in } 2^\circ, \text{ since } \int_0^1 \int_0^{2\pi} \|g(re^{i\theta})\| d\theta dr < \infty,$$

there exists  $u: [0, 1] \rightarrow ]0, \infty[$  continuous with  $u(r) \uparrow \infty$  as  $r \uparrow 1$  and

$$C := \int_0^1 u(r)r \int_0^{2\pi} \|g(re^{i\theta})\| d\theta dr < \infty. \text{ Then } w(r) := \frac{1}{u(r)}$$

$$f \in H^\infty(X): |\varphi(f)| = \left| \int_0^1 \int_0^{2\pi} \langle w(r)f(re^{i\theta}), u(r)g(re^{i\theta}) \rangle r d\theta dr \right| \leq C \|f\|_w.$$

3°  $\Rightarrow$  4°: With  $w$  as in 3°, the map  $f \mapsto wf$  is an isometric embedding ( $H^\infty(X)$ ,

$\|\cdot\|_w \hookrightarrow C_0(D; X) \subset C(\bar{D}; X)$  and  $\varphi$  is  $\|\cdot\|_w$ -continuous. By Singer's theorem [Si] (and Hahn-Banach), there exists  $m \in M(\bar{D}; X')$  vanishing on  $T$  (i.e.  $m \in M(D; X')$ ) with  $\varphi(f) = \iint_D \langle wf, dm \rangle = \iint_D \langle f, wdm \rangle, f \in H^\infty(X)$ .

$4^\circ \Rightarrow 1^\circ$ : For  $m \in M(D; X')$  representing  $\varphi$  as in  $4^\circ$ , the lemma (ii) yields  $\varphi(f) = \iint_D \langle f, dm \rangle = \int_T \langle f, B(m) \rangle d\lambda \forall f \in H^\infty(X) \subset H^\infty(X'')$ , where  $B[m] \in L^1(T; X')$ .

The proof of the equivalence of  $1^\circ-4^\circ$  is complete.

$2^\circ \Rightarrow 5^\circ$ : Bounded convergence.

$5^\circ \Rightarrow 3^\circ$ : Putting  $B := B_{H^\infty(X)}$ , condition  $5^\circ$  trivially implies that  $\varphi|B$  is sequentially continuous for  $\beta$ . The topology  $\kappa$  of uniform convergence on compact subsets of  $D$  is certainly weaker than  $\beta$ , and coincides with  $\beta$  on  $B$ , as is easy to see [RS, 3.7].  $\kappa$  being metrizable, the same holds for  $\beta|B$  so that  $\varphi|B$  is actually continuous for  $\beta$ . Now I apply Grothendieck's dual characterization of completeness [Sc, IV.6.2] for the locally convex space  $(H^\infty(X), \beta)$  and the topology of uniform convergence on norm-bounded sets (that is, on  $B$ ) on  $(H^\infty(X), \beta)'$ . By the equivalence of  $1^\circ$  and  $3^\circ$  already established, as a vector space  $(H^\infty(X), \beta)' = L^1(X')/H_0^1(X')$ , and the topology in question is given by the quotient norm, as proved below. In particular, this topology is complete, and Grothendieck's theorem yields that  $\varphi$  is continuous for  $\beta$ , i.e.  $3^\circ$ . It remains to establish the

CLAIM. For  $\Psi = g + H_0^1(X') \in L^1(X')/H_0^1(X')$  ( $g \in L^1(X')$ ),  $\|\Psi\|_{L^1(X')/H_0^1(X')} = \|\Psi\|_{L^1(X', X)/H_0^1(X')} = \|\Psi\|_{H^\infty(X)'}$ .

PROOF. i)  $\geq$  clear;  $\leq$ : Let  $\varepsilon > 0$  and  $h \in H_0^1(X')$  such that  $d(g, H_0^1(X')) \geq \|g - h\|_1 - \varepsilon$ . Now  $h_r := P_r * h \in H_0^1(X')$ , whence  $d(g, H_0^1(X')) \leq \overline{\lim}_{r \rightarrow 1} \|g - h_r\|_1 \leq$

$$\lim_{r \rightarrow 1} \|g - g_r\|_1 + \sup_{r < 1} \|g_r - h_r\|_1 = 0 + \|g - h\|_1 \leq d(g, H_0^1(X')) + \varepsilon.$$

ii)  $H^\infty(X)' = L^\infty(X)/H^\infty(X)^\perp = L^1(X', X)/H_0^1(X') \oplus_1 \dots$  by 1.2, 2.2 and the lemma on quotients of  $L$ -decompositions used already in the proof of 2.6.

3.6. REMARK. The theorem persists if  $H^\infty(X)$  is replaced by  $H^\infty(X)$  throughout.

3.7. REMARK. Every integral functional on  $H^\infty(X)$  is strongly integral if  $X'$  has RNP.

PROOF. The space of integral functionals  $L^1(X', X)/H_0^1(X')$  equals the space of strongly integral functionals  $L^1(X')/H_0^1(X')$  iff  $L^1(X', X) = L^1(X') + H_0^1(X')$ . Now if  $X'$  has RNP then  $L^1(X', X) = L^1(X')$  and the equality holds. The reverse implication is rather deep [LM, IV b] (take  $q = 1$  there).



#### §4. F. and M. Riesz Theorem for $H^\infty(B(H))$ .

Recall from 2.4, 2.6, 3.2 that for reflexive  $X$  the spaces  $H_0^1(X)$  and  $L^1(X)/H_0^1(X)$  are  $L$ -complemented in their biduals and have property (X). Now I establish the same for  $X = N(H)$ , the space of nuclear linear operators on separable Hilbert space. Since  $N(H)' = B(H)$  lacks RNP, the dual  $L^\infty(B, N)$  of  $L^1(N)$  does *not* coincide with  $L^\infty(B)$  [DU, IV.1.1] and the Yosida-Hewitt decomposition 1.2 is of no (direct) use. The strategy is rather to exploit the Takesaki  $L$ -decomposition [T 3, III.2.14] of the dual of the von Neumann algebra  $L^\infty(B, N)$ . In order to devolve this decomposition on the quotient  $(L^1(N)/H_0^1(N))' = L^\infty(B, N)/H^\infty(B)^\perp$  as in 2.6, an F. and M. Riesz theorem for  $H^\infty(B)$  is needed.

4.1. So let  $(H, (\cdot, \cdot))$  be a separable Hilbert space,  $B = B(H)$  resp.  $N = N(H)$  the space of bounded resp. nuclear (= trace class) operators on  $H$ . As is well-known,  $N$  is the (unique) predual of  $B$  under the dual pairing  $\langle a, b \rangle := \text{tr}(ab)$ ,  $a \in N$ ,  $b \in B$  [Pd, 3.4.13; T 3, III.3.9]. Therefore, by general vector-valued theory (1.1)  $L^1(N)' = L^\infty(B, N)$ , but in this operator situation the latter space has a more familiar description: Let  $L_w^\infty(B) := \{f: T \rightarrow B \text{ bounded: } \forall x, y \in H: (f(\cdot)x, y) \text{ measurable}\} / \{f = 0 \text{ a.e.}\}$ , equipped with the essential sup norm. Invoking the separability of  $H$  (hence of  $N(H)$ ) and [Pd, 4.6.11] it is routine to establish that  $L_w^\infty(B) = L^\infty(B, N)$  canonically. If an orthonormal basis  $(e_n)$  of  $H$  is fixed, then also  $L_w^\infty(B) = \{f: T \rightarrow B \text{ bounded: } \forall i, j: (f(\cdot)e_j, e_i) =: f_{ij} \in L^\infty\} / \{f = 0 \text{ a.e.}\}$ . In this description,  $H^\infty(B) = \{f \in L_w^\infty(B): \forall i, j: f_{ij} \in H^\infty\}$ . (Use that generally  $f \in L^\infty(X', X)$  is in  $H^\infty(X')$  iff  $\langle x, f \rangle \in H^\infty \forall x \in X$  and [Pd, 4.6.11] again.)

4.2. It is known that  $L_w^\infty(B)$  acts naturally as a von Neumann algebra on the Hilbert space  $L^2(H)$ ; these operators on  $L^2(H)$  are called decomposable [Pd, E 4.7.5; T 3, IV.7]. So  $L^1(N)$  is the unique predual of  $L_w^\infty(B)$ ; in fact,  $L^1(N)$  has property (X) [Go 2, V.4(4)].

4.3. *Takesaki decomposition* [T1; T3, III.2]: This is the noncommutative analogue of the Yosida-Hewitt decomposition. Let  $A$  be an arbitrary von Neumann algebra,  $A_*$  = predual of  $A$  = space of  $\sigma$ -weakly continuous (“normal”) functionals on  $A$ . There is a central projection  $z \in A'$  (identified with the universal enveloping von Neumann algebra of  $A$ ) such that  $A_* = A'_*z$ , where  $a'z \in A'$ ,  $(a'z)(a) := (az)(a')$  ( $a \in A$ ,  $a' \in A'$ ).  $z$  is called the support projection of  $A_* \subset A'$ . Then  $A_{\perp} := A'_*(1 - z)$  is the space of “singular” functionals on  $A$ .

**THEOREM** [T 3, III.2.14].  $A' = A_* \oplus_1 A_{\perp}$ .

The corresponding decomposition of  $\varphi \in A'$  is denoted  $\varphi = \varphi_* + \varphi_{\perp}$ :  $T: A' \rightarrow A_*$ ,  $T\varphi := \varphi_* = \varphi z$  is called Takesaki  $L$ -projection.

4.4. LEMMA. *Let  $p \in A$  be a projection,  $\varphi \in A'$ . The Takesaki decomposition of  $\varphi|_pAp$  is induced by that of  $\varphi$ .*

This lemma is analogous to 2.1. Note that  $pAp$  can be identified with a von Neumann algebra on  $\text{range}(p)$ .

PROOF. This is known [T 2, p. 365]. The argument is that if  $z \in A''$  is the support projection of  $A_* \subset A'$  then  $pz = zp = pzp \in pA''p = (pAp)''$  is the support projection of  $pA_*p = (pAp)_* \subset (pAp)'' = pA'p$ .

4.5. F. AND M. RIESZ THEOREM FOR  $H^\infty(B)$ : *Let  $B = B(H)$  as in 5.1,  $K = K(H) \subset B$  be the space of compact operators, and  $\varphi = \varphi_* + \varphi_\perp$  be the Takesaki decomposition (5.3) of a member  $\varphi \in L_w^\infty(B)'$ . If  $\varphi|_{H^\infty(B)} = 0$  then separately  $\varphi_*|_{H^\infty(B)} = 0 = \varphi_\perp|_{H^\infty(B)}$ . Moreover,  $\varphi_\perp$  also vanishes on  $C(K)$ .*

PROOF. First case:  $\dim H < \infty$ . Then of course  $L_w^\infty(B) = L^\infty(B)$ , and both the Takesaki (5.3) and the Yosida-Hewitt (2.2)  $L$ -projections in  $L^\infty(B)'$  have range  $L^1(N)$ . By uniqueness of the  $L$ -complement [HWW, 1.1.2], both decompositions coincide and the theorem is a special case of 2.2 resp. 2.3.

Second case:  $\dim H = \infty$ . The following finite-dimensional approximation is inspired by [Pi, 3.4]. Fix an orthonormal basis  $(e_n)_{n \in \mathbb{N}}$  of  $H$  and let  $p_n \in B$  be the orthogonal projection onto  $H_n := \text{lin}\{e_1, \dots, e_n\}$ . The space  $B_n := p_n B p_n$  consists of matrices with 0 outside the upper left  $n \times n$ -corner and can thus be identified with  $B(H_n)$ ; also,  $L^\infty(B_n) = p_n L_w^\infty(B) p_n$ , where  $p_n \in L_w^\infty(B)$  is considered now as the constant function  $p_n$  (these are the identifications after lemma 4.4). Let  $\varphi_n := \varphi|_{L^\infty(B_n)}$ , then  $\varphi|_{H^\infty(B)} = 0$  obviously implies  $\varphi_n|_{H^\infty(B_n)} = 0$ , hence  $\varphi_{n,*}|_{H^\infty(B_n)} = 0$  and  $\varphi_{n,\perp}|_{C(B_n)} = 0$  by the first case of the proof. Lemma 4.4 says  $\varphi_{n,*} = \varphi_{*,n} := \varphi_*|_{L^\infty(B_n)}$  and  $\varphi_{n,\perp} = \varphi_{\perp,n} := \varphi_\perp|_{L^\infty(B_n)}$ .

i) Let  $\varphi_* \in L_w^\infty(B)_*$  be given by  $g \in L^1(N)$ , and let  $f \in H^\infty(B)$ . In order to prove  $\varphi_*(f) = \int \langle g, f \rangle d\lambda = 0$ , consider  $0 = \varphi_*(p_n f(\cdot) p_n) = \int \langle g(t), p_n f(t) p_n \rangle \frac{dt}{2\pi}$ . Now pointwise  $p_n f(t) p_n \xrightarrow{n \rightarrow \infty} f(t)$  in the weak operator topology, hence [Pd, 4.6.14] for  $\sigma(B, N)$ . Thus  $\langle g(t), p_n f(t) p_n \rangle \xrightarrow{n \rightarrow \infty} \langle g(t), f(t) \rangle$  pointwise and dominated by  $\|f\|_{L_\bullet^\infty(B)} \|g(\cdot)\|_N \in L^1$ . By Lebesgue's theorem,  $\int \langle g, f \rangle d\lambda = \lim_{n \rightarrow \infty} \int \langle g, p_n f p_n \rangle d\lambda = 0$ .

ii) Now let  $f \in C(K)$ . In order to prove  $\varphi_\perp(f) = 0$  it suffices to observe that  $\lim_{n \rightarrow \infty} \|f(e^{it}) - p_n f(e^{it}) p_n\|_{K(H)} = 0$  uniformly in  $e^{it} \in T$ . Then norm continuity of  $\varphi_\perp$  suffices to conclude  $\varphi_\perp(f) = \lim_{n \rightarrow \infty} \varphi_\perp(p_n f p_n) = 0$ .

For the next corollary, note that  $N = K'$ , hence  $H_0^1(N) = H_0^1(N) = (C(K)/A(K))'$ .

4.6. COROLLARY 1.  $H_0^1(N)' = H_0^1(N) \oplus_1 (L_w^\infty(B)'_\perp \cap H^\infty(B)^\perp)$ .  $C(K)/A(K)$  is an  $M$ -ideal in its bidual  $L_w^\infty(B)/H^\infty(B)$ .

PROOF. The decomposition follows from theorems 4.3 and 4.5 (cf. 2.4). The second assertion follows from the last part of theorem 4.5:  $L_w^\infty(B)'_\perp \cap H^\infty(B)^\perp \subset C(K)^\perp$  (cf. 2.5).

REMARK. The  $M$ -embedded nature of  $C(K)/A(K)$  can also be derived from Page's theorem [Pa, Theorem 6], according to which  $C(K(H))/A(K(H))$  is isometrically isomorphic to the space of compact Hankel operators on the Hilbert space  $H^2(H)$ , cf. [HWW, III.1.8].

4.7. COROLLARY 2.  $L^1(N)/H_0^1(N)$  is  $L$ -embedded in its bidual.

PROOF. Theorems 4.3 and 4.5 (cf. 2.6).

As noted in section 2.6, it follows that  $L^1(N)/H_0^1(N)$  is weakly sequentially complete and even enjoys property (V\*). This points into the direction of a positive answer to a question of Pisier:  $H^\infty(B)$  is it a Grothendieck space? This question arises because both  $H^\infty$  and  $B$  are Grothendieck spaces and even have property (V) [Bo; Pf 2].

4.8. PROPOSITION.  $L^1(N)/H_0^1(N)$  has property (X).

PROOF. Similar to that of 4.5. Let  $\varphi = \varphi_* + \varphi_\perp \in L_w^\infty(B)'$  respect the  $w^*$ -convergence of  $wuC$  series in  $H^\infty(B)$ ; since  $\varphi_*$  does the same I can assume from the outset that  $\varphi = \varphi_\perp$  is singular and have to show that  $\varphi$  vanishes on  $H^\infty(B)$ . In the first case ( $\dim H < \infty$ ) this follows again by the argument "Takesaki = Yosida-Hewitt" and 3.2. For the second case ( $\dim H = \infty$ ) note first that  $\sum_{n=1}^\infty (p_n - p_{n-1})$  (where  $p_0 := 0$ ) is a  $wuC$  series in  $B$ , so that for every  $f \in H^\infty(B)$  both series  $\sum_{n=1}^\infty (p_n - p_{n-1})f$  and  $\sum_{n=1}^\infty f(p_n - p_{n-1})$  are  $wuC$  series in  $H^\infty(B)$  with  $w^*$ -limit  $f$ . Now fix  $f \in H^\infty(B)$ ; in order to prove  $\varphi(f) = 0$  it therefore suffices to prove  $\varphi(p_m f) = 0$  for every fixed  $m$ , and now it suffices to prove  $\varphi(p_m f p_n) = 0$  for every  $n \geq m$ . But  $p_m f p_n = p_n p_m f p_n \in H^\infty(B_n)$ ;  $\varphi|_{L^\infty(B_n)}$  is singular (4.4) and respects the  $w^*$ -convergence of  $wuC$  series in  $H^\infty(B_n)$ , hence vanishes on  $H^\infty(B_n)$  by the first case of the proof which is now complete.

REFERENCES

A T. Ando, *On the predual of  $H^\infty$* , Comment. Math., Tomus specialis in honorem L. Orlicz I, Warszawa, 1978.  
 BK K. Barbey, H. König, *Abstract Analytic Function Theory and Hardy Algebras*, Lecture Notes in Math. 593 (1977).

- BS K. D. Bierstedt, W. H. Summers, *Biduals of weighted Banach spaces of analytic functions*, J. Austral. Math. Soc. (Ser. A) 54 (1993), 70–79.
- Bo J. Bourgain,  $H^\infty$  is a Grothendieck space, *Studia Math.* 75 (1983), 193–216.
- Bu 1 A. V. Bukhvalov, *Hardy spaces of vector-valued functions*, Zap. Nauchn. Sem. LOMI 65 (1976), 5–16 (Russian); English translation; J. Soviet Math. 16 (1981), 1051–1059.
- Bu 2 A. V. Bukhvalov, *Optimization without compactness and its applications*. Unpublished manuscript, 1990.
- BD A. V. Bukhvalov, A. A. Danilevich, *Boundary properties of analytic and harmonic functions with values in Banach space*, Mat. Zametki 31 (1982), 203–214 (Russian); English translation: Math. Notes 31 (1982), 104–110.
- BL A. V. Bukhvalov, G. Ya. Lozanovskii, *On sets closed in measure in spaces of measurable function*. Trudy Moskov. Mat. Ob-va 34 (1977), 129–150 (Russian); English translation: Trans. Moscow Math. Soc. (1978), issue 2, 127–148.
- CG 1 M. Cambren, P. Greim, *The dual of a space of vector measures*, Math. Z. 180 (1982), 373–378.
- CG 2 M. Cambren, P. Greim, *Uniqueness of preduals for spaces of continuous vector functions*, Canad. Math. Bull. 31 (1988), 98–104.
- CV C. Castaing, M. Valadier, *Convex Analysis and Measurable Multifunctions*, Lecture Notes in Math. 580 (1977).
- Co J. B. Cooper, *Saks Spaces and Applications to Functional Analysis*, North Holland, Amsterdam etc. 1978 (North Holland Math. Studies 28, Notas de Matematica 64).
- DU J. Diestel, J. J. Uhl, Jr., *Vector Measures*, Math. Surveys 15 (1977).
- E G. A. Edgar, *An ordering for the Banach spaces*, Pacific J. Math. 108 (1983), 83–98.
- Ga J. B. Garnett, *Bounded Analytic Functions*, Academic Press, New York, 1981.
- Go 1 G. Godefroy, *Sous-espaces bien disposés de  $L^1$ -applications*, Trans. Amer. Math. Soc. 286 (1984), 227–249.
- Go 2 G. Godefroy, *Existence and uniqueness of isometric preduals: a survey*, Contemp. Math. 85 (1989), 131–193.
- GS G. Godefroy, P. Saab, *Quelques espaces de Banach ayant les propriétés (V) ou (V\*) de A. Pelczyński*, C. R. Acad. Sci. Paris 303 (1986), 503–506.
- GT G. Godefroy, M. Talagrand, *Classes d'espaces de Banach à prédual unique*, C.R. Acad. Sci. Paris 292 (1981), 323–325.
- HWW P. Harmand, D. Werner, W. Werner, *M-ideals in Banach Spaces and Banach Algebras*, Lecture Notes in Math. 1547 (1993).
- He 1 W. Hensgen, *Hardy-Räume vektorwertiger Funktionen*, Dissertation, München, 1986.
- He 2 W. Hensgen, *On weak sequential completeness of vector-valued  $L^1/H_0^1$* , Monografii matematice 47 (1992), Univ. Timisoara.
- He 3 W. Hensgen, *Contributions to the Geometry of Vector-Valued  $H^\infty$  and  $L^1/H_0^1$  Spaces*, Habilitationsschrift, Regensburg 1992.
- He 4 W. Hensgen, *Weak compactness in vector-valued  $L^1/H_0^1$* , Proc. Conf. on the Interaction between Functional Analysis, Harmonic Analysis and Probability, Columbia, Mo. 1994, LNPAM, Marcel Dekker.
- Ho K. Hoffmann, *Banach Spaces of Analytic Functions*, Prentice Hall, Inc., Englewood Cliffs (N.J.), 1962.
- Kh 1 V. P. Khavin, *Weak completeness of the space  $L^1/H_0^1$* , Vestnik Leningrad Univ. Math. 1973, no. 13, 77–81 (Russian); English translation: 6 (1979), 297–302.
- Kh 2 V. P. Khavin, *Spaces  $H^\infty$  and  $L^1/H_0^1$* , Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov 39 (1974), 120–148 (Russian); English translation: J. Sov. Math. 8 (1977), 86–108.
- LM C. Le Merdy, *Formes bilinéaires hankeliennes compactes sur  $H^2(X) \times H^2(Y)$* , Math. Ann. 293 (1992), 371–384.
- Pa L. B. Page, *Bounded and compact vectorial Hankel operators*, Trans. Amer. Math. Soc. 150 (1970), 529–539.
- Pd G. K. Pedersen, *Analysis Now*, Graduate Texts in Math. 118.
- Pe A. Pelczyński, *Banach Spaces of Analytic Functions and Absolutely Summing Operators*, AMS, Providence (Rh. I.) 1977 (Reg. Conf. Ser. in Math. 30).

- Pf 1 H. Pfützner, *L-summands in their bidual have Pelczyński's property ( $V^*$ )*, *Studia Math.* 104 (1993), 91–98.
- Pf 2 H. Pfützner, *Weak compactness in the dual of  $C^*$  algebras is determined commutatively*, *Math. Ann.* 298 (1994), 349–372.
- Pi G. Pisier, *Interpolation between  $H^p$  spaces and non-commutative generalizations*. *Pacific J. Math.* 155 (1992), 341–368.
- RR L. Rubel, J. V. Ryff, *The bounded weak\* topology and the bounded analytic functions*, *J. Funct. Anal.* 5 (1970), 167–183.
- RS L. A. Rubel, A. L. Shields, *The space of bounded analytic functions on a region*, *Ann. Inst. Fourier* 16 (1966), 235–277.
- Sc H. H. Schaefer, *Topological Vector Spaces*, Springer, New York, 4th printing 1980.
- Si I. Singer, *Sur les applications linéaires intégrales des espaces des fonctions continues*, *Rev. Roumaine Math. Pures Appl.* 4 (1959), 391–401.
- T 1 M. Takesaki, *On the conjugate space of operator algebra*, *Tôhoku Math. J.* 10 (1958), 194–203.
- T 2 M. Takesaki, *On the singularity of a positive linear functional on operator algebra*, *Proc. Japan Acad.* 35 (1959), 365–366.
- T 3 M. Takesaki, *Theory of Operator Algebras I*, Springer, New York, 1979.
- W P. Wojtaszczyk, *On projections in spaces of bounded analytic functions with applications*, *Studia Math.* 65 (1979), 147–173.

NWFI-MATHEMATIK

UNIVERSITÄT REGENSBURG

D-93040 REGENSBURG

e-mail address: wolfgang.hensgen@Mathematik.uni-regensburg.de