

ELEMENTARY NOTIONS OF LATTICE TRIGONOMETRY

OLEG KARPENKOV*

Introduction

0.1. *The goals of this paper and some background*

Consider a two-dimensional oriented real vector space and fix some full-rank lattice in it. A triangle or a polygon is said to be *lattice* if all its vertices belong to the lattice. The angles of any lattice triangle are said to be *lattice*.

In this paper we introduce and study *lattice trigonometric* functions of lattice angles. The lattice trigonometric functions are invariant under the action of the group of lattice-affine transformations (i.e. affine transformations preserving the lattice), like the ordinary trigonometric functions are invariant under the action of the group of Euclidean length preserving transformations of Euclidean space.

One of the initial goals of the present article is to make a complete description of lattice triangles up to the lattice-affine equivalence relation (see Theorem 2.2). The classification problem of convex lattice polygons becomes now classical. There is still no a good description of convex polygons. It is only known that the number of such polygons with lattice area bounded from above by n grows exponentially in $n^{1/3}$, while n tends to infinity (see the works of V. Arnold [2], and of I. Bárány and A. M. Vershik [3]).

We extend the geometric interpretation of ordinary continued fractions to define lattice sums of lattice angles and to establish relations on lattice tangents of lattice angles. Further, we describe lattice triangles in terms of *lattice sums* of lattice angles.

In present paper we also show a lattice version of the sine formula and introduce a relation between the lattice tangents for angles of lattice triangles and the numbers of lattice points on the edges of triangles (see Theorem 1.15).

* Partially supported by NWO-RFBR 047.011.2004.026 (RFBR 05-02-89000-NWO_a) grant, by RFBR SS-1972.2003.1 grant, by RFBR 05-01-02805-CNRS_La grant, and by RFBR grant 05-01-01012a.

Received October 17, 2006.

We conclude the paper with applications to toric varieties and some unsolved problems.

The study of lattice angles is an essential part of modern lattice geometry. Invariants of lattice angles are used in the study of lattice convex polygons and polytopes. Such polygons and polytopes play the principal role in Klein's theory of multidimensional continued fractions (see, for example, the works of F. Klein [14], V. I. Arnold [1], E. Korkina [16], M. Kontsevich and Yu. Suhov [15], G. Lachaud [17], and the author [10]).

Lattice polygons and polytopes of the lattice geometry are in the limelight of complex projective toric varieties (see for more information the works of V. I. Danilov [4], G. Ewald [5], T. Oda [18], and W. Fulton [6]). To illustrate, we deduce (in Appendix A) from Theorem 2.2 the corresponding global relations on the toric singularities for projective toric varieties associated to integer-lattice triangles. We also show the following simple fact: for any collection with multiplicities of complex-two-dimensional toric algebraic singularities there exists a complex-two-dimensional toric projective variety with the given collection of toric singularities (this result seems to be classical, but it is missing in the literature).

The studies of lattice angles and measures related to them were started by A. G. Khovanskii, A. Pukhlikov in [12] and [13] in 1992. They introduced and investigated special additive polynomial measure for the extended notion of polytopes. The relations between sum-formulas of lattice trigonometric functions and lattice angles in Khovanskii-Pukhlikov sense are unknown to the author.

0.2. Some distinctions between lattice and Euclidean cases

Lattice trigonometric functions and Euclidean trigonometric functions have much in common. For example, the values of lattice tangents and Euclidean tangents coincide in a special natural system of coordinates. Nevertheless, lattice geometry differs a lot from Euclidean geometry. We show this with the following four examples.

1. The angles $\angle ABC$ and $\angle CBA$ are always congruent in Euclidean geometry, but not necessary lattice-congruent in lattice geometry.

2. In Euclidean geometry for any $n \geq 3$ there exist a regular polygon with n vertices, and any two regular polygons with the same number of vertices are homothetic to each other. In lattice geometry there are only six non-homothetic regular lattice polygons: two triangles (distinguished by lattice tangents of angles), two quadrangles, and two octagons. (See a more detailed description in [11].)

3. In Appendix B we will consider three natural criteria for triangle congruence in Euclidean geometry. Only the first criterion can be taken to the case of

lattice geometry. The others two are false in lattice trigonometry. (We refer to Appendix B.)

4. There exist two non-congruent right angles in lattice geometry. (See Corollary 1.12.)

0.3. Description of the paper

This paper is organized as follows.

We start in Section 1 with some general notation of lattice geometry. We define ordinary lattice angles, and the functions lattice sine, tangent, and cosine on the set of ordinary lattice angles, and lattice arctangent for rationals greater than or equal 1. Further we indicate their basic properties. We proceed with the geometrical interpretation of lattice tangents in terms of ordinary continued fractions. In conclusion of Section 1 we study the basic properties of angles in lattice triangles.

In Section 2 we introduce the sum formula for the lattice tangents of ordinary lattice angles of lattice triangles. The sum formula is a lattice generalization of the following Euclidean statement: three angles are the angles of some triangle iff their sum equals π .

Further in Section 3 we introduce the notion of extended lattice angles and their normal forms and give the definition of sums of extended and ordinary lattice angles. Here we extend the notion of sails in the sense of Klein: we define and study oriented broken lines at unit distance from lattice points.

In Section 4 we finally prove the first statement of the theorem on sums of lattice tangents for angles in lattice triangles. In this section we also describe some relations between continued fractions for lattice oriented broken lines and the lattice tangents for the corresponding extended lattice angles. Further we give a necessary and sufficient condition for an ordered n -tuple of angles to be the angles of some convex lattice polygon.

We conclude this paper with three appendices. In Appendix A we describe applications to theory of complex projective toric varieties mentioned above. Further in Appendix B we formulate criterions of lattice congruence for lattice triangles. Finally in Appendix C we give a list of unsolved problems and questions.

ACKNOWLEDGEMENT. The author is grateful to V. I. Arnold for constant attention to this work, I. Bárány, A. G. Khovanskii, V. M. Kharlamov, J.-M. Kantor, D. Zvonkine, and D. Panov for useful remarks and discussions, and Université Paris-Dauphine — CEREMADE for the hospitality and excellent working conditions.

1. Definitions and elementary properties of lattice trigonometric functions

1.1. Preliminary notions and definitions

By $\gcd(n_1, \dots, n_k)$ and by $\text{lcm}(n_1, \dots, n_k)$ we denote the greater common divisor and the less common multiple of the nonzero integers n_1, \dots, n_k respectively. Suppose that a, b be arbitrary integers, and c be an arbitrary positive integer. We write that $a \equiv b \pmod{c}$ if the remainders of a and b modulo c coincide.

1.1.1. Lattice notation. Here we define the main objects of lattice geometry, their lattice characteristics, and the relation of \mathcal{L} -congruence (lattice-congruence).

Consider \mathbb{R}^2 and fix some orientation and some lattice in it. A straight line is said to be *lattice* if it contains at least two distinct lattice points. A ray is said to be *lattice* if its vertex is a lattice point, and it contains lattice points distinct from its vertex. An angle (i.e. the union of two rays with the common vertex) is said to be *ordinary lattice* (or just *ordinary* for short) if the rays defining it are lattice. A segment is called *lattice* if its endpoints are lattice points.

By a *convex polygon* we mean a convex hulls of a finite number of points that do not lie in a straight line. A straight line π is said to be *supporting* a convex polygon P , if the intersections of P and π is not empty, and the whole polygon P is contained in one of the closed half-planes bounded by π . An intersection of a polygon P with its supporting straight line is called a *vertex* or an *edge* of the polygon if the dimension of intersection is zero, or one respectively.

A triangle (or convex polygon) is said to be *lattice* if all its vertices are lattice points. A lattice triangle is said to be *simple* if the vectors corresponding to its edges generate the lattice.

The affine transformation is called *\mathcal{L} -affine* if it preserves the set of all lattice points. Consider two arbitrary (not necessary lattice in the above sense) sets. We say that these two sets are *\mathcal{L} -congruent* to each other if there exist a *\mathcal{L} -affine* transformation of \mathbb{R}^2 taking the first set to the second.

DEFINITION 1.1. The *lattice length* of a lattice segment AB is the ratio between the Euclidean length of AB and the length of the basic lattice vector for the straight line containing this segment. We denote the lattice length by $\ell(AB)$.

By the (non-oriented) *lattice area* of the convex polygon P we will call the ratio of the Euclidean area of the polygon and the area of any lattice simple triangle, and denote it by $\text{IS}(P)$.

Two lattice segments are \mathcal{L} -congruent iff they have equal lattice lengths. The lattice area of the convex polygon is well-defined and is proportional to the Euclidean area of the polygon.

1.1.2. Finite ordinary continued fractions. For any finite sequence (a_0, a_1, \dots, a_n) where the elements a_1, \dots, a_n are positive integers and a_0 is an arbitrary integer we associate the following rational number q :

$$q = a_0 + \frac{1}{a_1 + \frac{1}{\ddots + \frac{1}{a_{n-1} + \frac{1}{a_n}}}}$$

This representation of the rational q is called an *ordinary continued fraction* for q and denoted by $[a_0, a_1, \dots, a_n]$.

An ordinary continued fraction $[a_0, a_1, \dots, a_n]$ is said to be *odd* if $n + 1$ is odd, and *even* if $n + 1$ is even. Note that if $a_n \neq 1$ then $[a_0, a_1, \dots, a_n] = [a_0, a_1, \dots, a_n - 1, 1]$. Let us formulate the following classical theorem.

THEOREM 1.2. *For any rational there exist exactly one odd ordinary continued fraction and exactly one even ordinary continued fraction.*

1.2. Definition of lattice trigonometric functions

In this subsection we define the functions lattice sine, tangent, and cosine on the set of ordinary lattice angles and formulate their basic properties. We describe a geometric interpretation of lattice trigonometric functions in terms of ordinary continued fractions. Then we give the definitions of ordinary angles that are adjacent, transpose, and opposite interior to the given angles. We use the notions of adjacent and transpose ordinary angles to define ordinary lattice right angles.

Let $A, O,$ and B be three lattice points that do not lie in the same straight line. We denote the ordinary angle with the vertex at O and the rays OA and OB by $\angle AOB$.

One can chose any other lattice point C in the open lattice ray OA and any lattice point D in the open lattice ray OB . For us the angle $\angle AOB$ coincides with $\angle COD$. We denote this by $\angle AOB = \angle COD$.

DEFINITION 1.3. Two ordinary angles $\angle AOB$ and $\angle A'O'B'$ are said to be \mathcal{L} -congruent if there exist a \mathcal{L} -affine transformation that takes the point O to O' and the rays OA and OB to the rays $O'A'$ and $O'B'$ respectively. We denote this as follows: $\angle AOB \cong \angle A'O'B'$.

Here we note that the relation $\angle AOB \cong \angle BOA$ holds only for special ordinary angles. (See also below in Subsubsection 1.2.4.)

1.2.1. Definition of lattice sine, tangent, and cosine for an ordinary lattice angle. Consider an arbitrary ordinary angle $\angle AOB$. Let us associate a special basis to this angle. Denote by \bar{v}_1 and by \bar{v}_2 the lattice vectors generating the rays of the angle:

$$\bar{v}_1 = \frac{\overline{OA}}{l\ell(OA)}, \quad \text{and} \quad \bar{v}_2 = \frac{\overline{OB}}{l\ell(OB)}.$$

The set of lattice points at unit lattice distance from the lattice straight line OA coincides with the set of all lattice points of two lattice straight lines parallel to OA . Since the vectors \bar{v}_1 and \bar{v}_2 are linearly independent, the ray OB intersects exactly one of the above two lattice straight lines. Denote this straight line by l . The intersection point of the ray OB with the straight line l divides l into two parts. Choose one of the parts which lies in the complement to the convex hull of the union of the rays OA and OB , and denote by D the lattice point closest to the intersection of the ray OB with the straight line l (see Figure 1).


Now we choose the vectors $\bar{e}_1 = \bar{v}_1$ and $\bar{e}_2 = \overline{OD}$. These two vectors are linearly independent and generate the lattice. The basis (\bar{e}_1, \bar{e}_2) is said to be *associated* to the angle $\angle AOB$.

Since (\bar{e}_1, \bar{e}_2) is a basis, the vector \bar{v}_2 has a unique representation of the form:

$$\bar{v}_2 = x_1\bar{e}_1 + x_2\bar{e}_2,$$

where x_1 and x_2 are some integers.

DEFINITION 1.4. In the above notation, the coordinates x_2 and x_1 are said to be the *lattice sine* and the *lattice cosine* of the ordinary angle $\angle AOB$ respectively. The ratio of the lattice sine and the lattice cosine (x_2/x_1) is said to be the *lattice tangent* of $\angle AOB$.



$$\begin{aligned} \bar{v}_2 &= 5\bar{e}_1 + 7\bar{e}_2 \\ l\sin \angle AOB &= 7 \\ l\cos \angle AOB &= 5 \\ l\tan \angle AOB &= 7/5 \end{aligned}$$

FIGURE 1. An ordinary angle $\angle AOB$ and its lattice trigonometric functions.

Figure 1 shows an example of lattice angle with the lattice sine equals 7 and the lattice cosine equals 5.

Let us briefly enumerate some elementary properties of lattice trigonometric functions.

PROPOSITION 1.5. a) *The lattice sine and cosine of any ordinary angle are relatively-prime positive integers.*

b) *The values of lattice trigonometric functions for \mathcal{L} -congruent ordinary angles coincide.*

c) *The lattice sine of an ordinary angle coincide with the index of the sublattice generated by all lattice vectors of two angle rays in the lattice.*

d) *For any ordinary angle α the following inequalities hold:*

$$l\sin \alpha \geq l\cos \alpha, \quad \text{and} \quad l\tan \alpha \geq 1.$$

The equalities hold iff the lattice vectors of the angle rays generate the whole lattice.

e) (Description of lattice angles) *Two ordinary angles α and β are \mathcal{L} -congruent iff $l\tan \alpha = l\tan \beta$.*

1.2.2. *Lattice arctangent.* Let us fix the origin O and a lattice basis \bar{e}_1 and \bar{e}_2 .

DEFINITION 1.6. Consider an arbitrary rational $p \geq 1$. Let $p = m/n$, where m and n are positive integers. Suppose $A = O + \bar{e}_1$, and $B = O + n\bar{e}_1 + m\bar{e}_2$. The ordinary angle $\angle AOB$ is said to be the *arctangent of p in the fixed basis* and denoted by $larctan(p)$.

The invariance of lattice tangents immediately implies the following properties.

PROPOSITION 1.7. a) *For any rational $s \geq 1$, we have: $l\tan(larctan s) = s$.*

b) *For any ordinary angle α the following holds: $larctan(l\tan \alpha) \cong \alpha$.*

1.2.3. *Lattice tangents, length-sine sequences, sails, and continued fractions.* Let us start with the notion of sails for the ordinary angles. This notion is taken from theory of multidimensional continued fractions in the sense of Klein (see, for example, the works of F. Klein [14], and V. Arnold [1]).

Consider an ordinary angle $\angle AOB$. Let also the vectors \overline{OA} and \overline{OB} be linearly independent, and of unit lattice length. Denote the closed convex solid cone for the ordinary angle $\angle AOB$ by $C(AOB)$. The boundary of the convex hull of all lattice points of the cone $C(AOB)$ except the origin is homeomorphic to the straight line. This boundary contains the points A and B . The closed part of this boundary contained between the points A and B is called the *sail* for the cone $C(AOB)$.

A lattice point of the sail is said to be a *vertex* of the sail if there is no lattice segment of the sail containing this point in the interior. The sail of the cone $C(AOB)$ is a broken line with a finite number of vertices and without self intersections. Let us orient the sail in the direction from A to B , and denote the vertices of the sail by V_i (for $0 \leq i \leq n$) according to the orientation of the sail (such that $V_0 = A$, and $V_n = B$).

DEFINITION 1.8. Let the vectors \overline{OA} and \overline{OB} of the ordinary angle $\angle AOB$ be linearly independent, and of unit lattice length. Let V_i , where $0 \leq i \leq n$, be the vertices of the corresponding sail. The sequence of lattice lengths and sines

$$(\ell(V_0V_1), \text{lsin } \angle V_0V_1V_2, \ell(V_1V_2), \text{lsin } \angle V_1V_2V_3, \dots, \ell(V_{n-2}V_{n-1}), \text{lsin } \angle V_{n-2}V_{n-1}V_n, \ell(V_{n-1}V_n))$$

is called the *lattice length-sine sequence* for the ordinary angle $\angle AOB$. Further we say *LLS-sequence* for short.

REMARK 1.9. The elements of the lattice LLS-sequence for any ordinary angle are positive integers. The LLS-sequences of \mathcal{L} -congruent ordinary angles coincide.

THEOREM 1.10. Let $(a_0, a_1, \dots, a_{2n-3}, a_{2n-2})$ be the LLS-sequence for the ordinary angle $\angle AOB$. Then the lattice tangent of the ordinary angle $\angle AOB$ equals to the value of the following ordinary continued fraction

$$[a_0, a_1, \dots, a_{2n-3}, a_{2n-2}].$$

On Figure 2 we show an example of an ordinary angle with tangent equivalent to $7/5$.




FIGURE 2. $\text{ltan } \angle AOB = \frac{7}{5} = 1 + \frac{1}{2+1/2}$.

Further in Theorem 3.5 we formulate and prove a general statement for generalized sails and signed lattice length-sine sequences. In the proof of Theorem 3.5 we refer only on the preceding statements and definitions of Subsection 3.1,

that are independent of the statements and theorems of all previous sections. For these reasons we skip now the proof of Theorem 1.10 (see also Remark 3.6).

1.2.4. Adjacent, transpose, and opposite interior ordinary angles. An ordinary angle $\angle BOA$ is said to be *transpose* to the ordinary angle $\angle AOB$. We denote it by $(\angle AOB)^t$. An ordinary angle $\angle BOA'$ is said to be *adjacent* to an ordinary angle $\angle AOB$ if the points A , O , and A' are contained in the same straight line, and the point O lies between A and A' . We denote the ordinary angle $\angle BOA'$ by $\pi - \angle AOB$. The ordinary angle is said to be *right* if it is \mathcal{L} -congruent to the adjacent and to the transpose ordinary angles.

It immediately follows from the definition, that for any ordinary angle α the angles $(\alpha^t)^t$ and $\pi - (\pi - \alpha)$ are \mathcal{L} -congruent to α .

In the next theorem we use the following notion. Suppose that some integers a, b and c , where $c \geq 1$, satisfy the following: $ab \equiv 1 \pmod{c}$. Then we denote $a \equiv (b \pmod{c})^{-1}$.

THEOREM 1.11. *Consider an ordinary angle α . If $\alpha \cong \text{larctan}(1)$, then*

$$\alpha^t \cong \pi - \alpha \cong \text{larctan}(1).$$

Suppose now, that $\alpha \not\cong \text{larctan}(1)$, then

$$\begin{aligned} \text{l sin}(\alpha^t) &= \text{l sin} \alpha, & \text{l cos}(\alpha^t) &\equiv (\text{l cos} \alpha \pmod{\text{l sin} \alpha})^{-1}; \\ \text{l sin}(\pi - \alpha) &= \text{l sin} \alpha, & \text{l cos}(\pi - \alpha) &\equiv (-\text{l cos} \alpha \pmod{\text{l sin} \alpha})^{-1}. \end{aligned}$$

Note also, that $\pi - \alpha \cong \text{larctan}^t\left(\frac{\text{l tan} \alpha}{\text{l tan}(\alpha) - 1}\right)$.

Theorem 1.11 (after applying Theorem 1.10) immediately reduces to the theorem of P. Popescu-Pampu. We refer the readers to his work [19] for the proofs.

1.2.5. Right ordinary lattice angles. It turns out that in lattice geometry there exist exactly two lattice non-equivalent right ordinary angles.

COROLLARY 1.12. *Any ordinary right angle is \mathcal{L} -congruent to exactly one of the following two angles: $\text{larctan}(1)$, or $\text{larctan}(2)$.*

Consider two lattice parallel distinct straight lines AB and CD , where A , B , C , and D are lattice points. Let the points A and D be in different open half-planes with respect to the straight line BC . Then the ordinary angle $\angle ABC$ is called *opposite interior* to the ordinary angle $\angle DCB$. Further we use the following proposition on opposite interior ordinary angles.

PROPOSITION 1.13. *Two opposite interior to each other ordinary angles are \mathcal{L} -congruent.*

The proof is left for the reader as an exercise.

1.3. Basic lattice trigonometry of lattice angles in lattice triangles

In this subsection we introduce the sine formula for angles and edges of lattice triangles. Further we show how to find the lattice tangents of all angles and the lattice lengths of all edges of any lattice triangle, if the lattice lengths of two edges and the lattice tangent of the angle between them are given.

Let A, B, C be three distinct and not collinear lattice points. We denote the lattice triangle with the vertices A, B , and C by $\triangle ABC$. The lattice triangles $\triangle ABC$ and $\triangle A'B'C'$ are said to be \mathcal{L} -congruent if there exist a \mathcal{L} -affine transformation which takes the point A to A' , B to B' , and C to C' respectively. We denote: $\triangle ABC \cong \triangle A'B'C'$.

PROPOSITION 1.14 (The sine formula for lattice triangles). *The following holds for any lattice triangle $\triangle ABC$.*

$$\frac{\ell(AB)}{\text{lsin } \angle BCA} = \frac{\ell(BC)}{\text{lsin } \angle CAB} = \frac{\ell(CA)}{\text{lsin } \angle ABC} = \frac{\ell(AB)\ell(BC)\ell(CA)}{\text{IS}(\triangle ABC)}.$$

PROOF. The statement of Proposition 1.14 follows directly from the definition of lattice sine.

Suppose that we know the lattice lengths of the edges AB, AC and the lattice tangent of $\angle BAC$ in the triangle $\triangle ABC$. Now we show how to restore the lattice length and the lattice tangents for the the remaining edge and ordinary angles of the triangle.

For the simplicity we fix some lattice basis and use the system of coordinates OXY corresponding to this basis (denoted $(*, *)$).

THEOREM 1.15. *Consider some triangle $\triangle ABC$. Let*

$$\ell(AB) = c, \quad \ell(AC) = b, \quad \text{and} \quad \angle CAB \cong \alpha.$$

Then the ordinary angles $\angle BCA$ and $\angle ABC$ are defined in the following way.

$$\angle BCA \cong \begin{cases} \pi - \text{larctan}\left(\frac{c \text{lsin } \alpha}{c \text{lcos } \alpha - b}\right) & \text{if } c \text{lcos } \alpha > b \\ \text{larctan}(1) & \text{if } c \text{lcos } \alpha = b \\ \text{larctan}^t\left(\frac{c \text{lsin } \alpha}{b - c \text{lcos } \alpha}\right) & \text{if } c \text{lcos } \alpha < b, \end{cases}$$

$$\angle ABC \cong \begin{cases} \left(\pi - \text{larctan}\left(\frac{b \text{lsin}(\alpha')}{b \text{lcos}(\alpha') - c}\right)\right)^t & \text{if } b \text{lcos}(\alpha') > c \\ \text{larctan}(1) & \text{if } b \text{lcos}(\alpha') = c \\ \text{larctan}\left(\frac{b \text{lsin}(\alpha')}{c - b \text{lcos}(\alpha')}\right) & \text{if } b \text{lcos}(\alpha') < c. \end{cases}$$

For the lattice length of the edge CB we have

$$\frac{l(CB)}{l \sin \alpha} = \frac{b}{l \sin \angle ABC} = \frac{c}{l \sin \angle BCA}.$$

PROOF. Let $\alpha \cong \text{larctan}(p/q)$, where $\text{gcd}(p, q) = 1$. Then $\triangle CAB \cong \triangle DOE$ where $D = (b, 0)$, $O = (0, 0)$, and $E = (qc, pc)$. Let us now find the ordinary angle $\angle EDO$. Denote by Q the point $(qc, 0)$. If $qc - b = 0$, then $\angle BCA = \angle EDO = \text{larctan } 1$. If $qc - b \neq 0$, then we have

$$\angle QDE \cong \text{larctan} \left(\frac{cp}{|cq - b|} \right) \cong \text{larctan} \left(\frac{c \, l \sin \alpha}{|c \, l \cos \alpha - b|} \right).$$

The expression for $\angle BCA$ follows directly from the above expression for $\angle QDE$, since $\angle BCA \cong \angle QDE$. (See Figure 3: here $l(OD) = b$, $l(OQ) = c \, l \cos \alpha$, and therefore $l(DQ) = |c \, l \cos \alpha - b|$.)




FIGURE 3. Three possible configuration of points O , D , and Q .

To obtain the expression for $\angle ABC$ we consider the triangle $\triangle BAC$. Calculate $\angle CBA$ and then transpose all ordinary angles in the expression. Since

$$\begin{aligned} lS(ABC) &= l(AB) \, l(AC) \, l \sin \angle CAB \\ &= l(BA) \, l(BC) \, l \sin \angle BCA \\ &= l(CB) \, l(CA) \, l \sin \angle ABC, \end{aligned}$$

we have the last statement of the theorem.

2. Theorem on sum of lattice tangents for the ordinary lattice angles of lattice triangles. Proof of its second statement

Throughout this section we fix some lattice basis and use the system of coordinates OXY corresponding to this basis.

2.1. Finite continued fractions with not necessary positive elements

We start this section with the notation for finite continued fractions with not necessary positive elements. Let us extend the set of rationals \mathbb{Q} with the operations $+$ and $1/*$ on it with the element ∞ . We pose $q \pm \infty = \infty$, $1/0 = \infty$, $1/\infty = 0$ (we do not define $\infty \pm \infty$ here). Denote this extension by $\overline{\mathbb{Q}}$.

For any finite sequence of integers (a_0, a_1, \dots, a_n) we associate an element q of $\overline{\mathbb{Q}}$:

$$q = a_0 + \frac{1}{a_1 + \frac{1}{\ddots + \frac{1}{a_{n-1} + \frac{1}{a_n}}}}$$

and denote it by $]a_0, a_1, \dots, a_n[$.

Let q_i be some rationals, $i = 1, \dots, k$. Suppose that the odd continued fraction for q_i is $]a_{i,0}, a_{i,1}, \dots, a_{i,2n_i}[$ for $i = 1, \dots, k$. We denote by $]q_1, q_2, \dots, q_n[$ the following number

$$]a_{1,0}, a_{1,1}, \dots, a_{1,2n_1}, a_{2,0}, a_{2,1}, \dots, a_{2,2n_2}, \dots, a_{k,0}, a_{k,1}, \dots, a_{k,2n_k}[$$

2.2. Formulation of the theorem and proof of its second statement

In Euclidean geometry the sum of Euclidean angles of the triangle equals π . For any 3-tuple of angles with the sum equals π there exist a triangle with these angles. Two Euclidean triangles with the same angles are homothetic. Let us show a generalization of these statements to the case of lattice geometry.

Let n be an arbitrary positive integer, and $A = (x, y)$ be an arbitrary lattice point. Denote by nA the point (nx, ny) .

DEFINITION 2.1. Consider any convex polygon or broken line with vertices A_0, \dots, A_k . The polygon or broken line $nA_0 \dots nA_k$ is called n -multiple (or multiple) to the given polygon or broken line.

THEOREM 2.2 (On sum of lattice tangents of angles in lattice triangles).

a) Let $(\alpha_1, \alpha_2, \alpha_3)$ be an ordered 3-tuple of ordinary angles. There exists a triangle with three consecutive ordinary angles \mathcal{L} -congruent to α_1, α_2 , and

α_3 iff there exists $i \in \{1, 2, 3\}$ such that the angles $\alpha = \alpha_i$, $\beta = \alpha_{i+1 \pmod{3}}$, and $\gamma = \alpha_{i+2 \pmod{3}}$ satisfy the following conditions:

- i) for $A =]\tan \alpha, -1, \tan \beta[$ the following holds $A < 0$, or $A > \tan \alpha$, or $A = \infty$;
- ii) $]\tan \alpha, -1, \tan \beta, -1, \tan \gamma[= 0$.

b) Let the consecutive ordinary angles of some triangle be α , β , and γ . Then this triangle is multiple to the triangle with vertices $A_0 = (0, 0)$, $B_0 = (\lambda_2 \operatorname{lcos} \alpha, \lambda_2 \operatorname{lsin} \alpha)$, and $C_0 = (\lambda_1, 0)$, where

$$\lambda_1 = \frac{\operatorname{lcm}(\operatorname{lsin} \alpha, \operatorname{lsin} \beta, \operatorname{lsin} \gamma)}{\operatorname{gcd}(\operatorname{lsin} \alpha, \operatorname{lsin} \gamma)}, \quad \text{and} \quad \lambda_2 = \frac{\operatorname{lcm}(\operatorname{lsin} \alpha, \operatorname{lsin} \beta, \operatorname{lsin} \gamma)}{\operatorname{gcd}(\operatorname{lsin} \alpha, \operatorname{lsin} \beta)}.$$

Let us say a few words about the essence of the theorem. In Euclidean geometry on the plane the condition on the angles of triangles can be rewritten with tangent functions in the following way. A triangle with angles exists α , β , and γ iff $\tan(\alpha + \beta + \gamma) = 0$ and $\tan(\alpha + \beta) \notin [0; \tan \alpha]$ (here without lose of generality we suppose that α is acute). Theorem 2.2 is a translation of this condition into lattice case.

In addition we say that there is no good description of lattice polygons terms of lattice invariants at present. Theorem 2.2 gives such description for the case of triangles.

At this moment we do not have the necessary notation to prove the first statement of Theorem 2.2. For a proof we need first to define extended angles and their sums, and study their properties. We give a proof further in Subsections 4.2 and 4.3. We prove the second statement of the theorem below in this subsection.

REMARK 2.3. Note that the statement of Theorem 2.2a holds only for odd continued fractions for the tangents of the correspondent angles. We illustrate this with the following example. Consider a lattice triangle with the lattice area equals 7 and all angles \mathcal{L} -congruent to $\operatorname{larctan} 7/3$. If we take the odd continued fractions $7/3 = [2, 2, 1]$ for all lattice angles of the triangle, then we have

$$]2, 2, 1, -1, 2, 2, 1, -1, 2, 2, 1[= 0.$$

If we take the even continued fractions $7/3 = [2, 3]$ for all angles of the triangle, then we have

$$]2, 3, -1, 2, 3, -1, 2, 3[= \frac{35}{13} \neq 0.$$

PROOF OF THE SECOND STATEMENT OF THEOREM 2.2. Consider a triangle $\triangle ABC$ with ordinary angles α , β , and γ (at vertices at A , B , and C respectively). Suppose that for any $k > 1$ and any lattice triangle $\triangle KLM$ the triangle $\triangle ABC$ is not \mathcal{L} -congruent to the k -multiple of $\triangle KLM$. In other words, we have

$$\gcd(\ell(AB), \ell(BC), \ell(CA)) = 1.$$

Suppose that S is the lattice area of $\triangle ABC$. Then by the sine formula the following holds

$$\begin{cases} \ell(AB)\ell(AC) = S/\sin \alpha \\ \ell(BC)\ell(BA) = S/\sin \beta \\ \ell(CA)\ell(CB) = S/\sin \gamma \end{cases}.$$

Since $\gcd(\ell(AB), \ell(BC), \ell(CA)) = 1$, we have $\ell(AB) = \lambda_1$ and $\ell(AC) = \lambda_2$.

Therefore, the lattice triangle $\triangle ABC$ is \mathcal{L} -congruent to the lattice triangle $\triangle A_0B_0C_0$ of the theorem.

3. Extension of ordinary lattice angles. Notion of sums of lattice angles

Throughout this section we work in with an oriented two-dimensional real vector space and a fixed lattice in it. We again fix some (positively oriented) lattice basis and use the system of coordinates OXY corresponding to this basis.

The \mathcal{L} -affine transformation is said to be *proper* if it is orientation-preserving (we denote it by \mathcal{L}_+ -affine transformation).

We say that two sets are \mathcal{L}_+ -congruent to each other if there exist a \mathcal{L}_+ -affine transformation of \mathbb{R}^2 taking the first set to the second.

3.1. On a particular generalization of sails in the sense of Klein

In this subsection we introduce the definition of an oriented broken lines at unit lattice distance from a lattice point. This notion is a direct generalization of the notion of a sail in the sense of Klein (see page 167 for the definition of a sail). We extend the definition of LLS-sequences and continued fractions to the case of these broken lines. We show that extended LLS-sequence for oriented broken lines uniquely identifies the \mathcal{L}_+ -congruence class of the corresponding broken line. Further, we study the geometrical interpretation of the corresponding continued fraction.

3.1.1. Definition of a lattice signed length-sine sequence. Let us extend the definition of LLS-sequence to the case of certain broken lines.

For the 3-tuples of lattice points $A, B,$ and C we define the function sgn as follows:

$$\text{sgn}(ABC) = \begin{cases} +1, & \text{if the pair of vectors } \overline{BA} \text{ and } \overline{BC} \text{ defines the positive orientation.} \\ 0, & \text{if the points } A, B, \text{ and } C \text{ are contained in the same straight line.} \\ -1, & \text{if the pair of vectors } \overline{BA} \text{ and } \overline{BC} \text{ defines the negative orientation.} \end{cases}$$

We also denote by $\text{sign} : \mathbb{R} \rightarrow \{-1, 0, 1\}$ the sign function over reals.

A segment AB is said to be *at unit distance* from the point C if the lattice vectors of the segment AB , and the vector \overline{AC} generate the lattice.

A union of (ordered) lattice segments $A_0A_1, A_1A_2, \dots, A_{n-1}A_n$ ($n > 0$) is said to be a *lattice oriented broken line* and denoted by $A_0A_1A_2 \dots A_n$ if any two consecutive segments are not contained in the same straight line. We also say that the lattice oriented broken line $A_nA_{n-1}A_{n-2} \dots A_0$ is *inverse* to the lattice oriented broken line $A_0A_1A_2 \dots A_n$.

DEFINITION 3.1. Consider a lattice oriented broken line and a lattice point V in the complement to this line. The broken line is said to be *at unit distance from the point V* (or *V -broken line for short*) if all edges of the broken line are at unit distance from V .

Let us now associate to any lattice oriented V -broken line for some lattice point V the following sequence of non-zero elements.

DEFINITION 3.2. Let $A_0A_1 \dots A_n$ be a lattice oriented V -broken line. The sequence of integers (a_0, \dots, a_{2n-2}) defined as follows:

$$\begin{aligned} a_0 &= \text{sgn}(A_0VA_1) \ell(A_0A_1), \\ a_1 &= \text{sgn}(A_0VA_1) \text{sgn}(A_1VA_2) \text{sgn}(A_0A_1A_2) \text{lsin } \angle A_0A_1A_2, \\ a_2 &= \text{sgn}(A_1VA_2) \ell(A_1A_2), \\ &\dots \\ a_{2n-3} &= \text{sgn}(A_{n-2}VA_{n-1}) \text{sgn}(A_{n-1}VA_n) \\ &\quad \text{sgn}(A_{n-2}A_{n-1}A_n) \text{lsin } \angle A_{n-2}A_{n-1}A_n, \\ a_{2n-2} &= \text{sgn}(A_{n-1}VA_n) \ell(A_{n-1}A_n), \end{aligned}$$

is called an *lattice signed length-sine sequence* for the lattice oriented V -broken line. Further we will say *LSSLS-sequence* for short.

The element $]a_0, a_1, \dots, a_{2n-2}[$ of $\overline{\mathbb{Q}}$ is called the *continued fraction for the broken line $A_0A_1 \dots A_n$* .

If we take LSLS-sequence for some broken line which is a sail, than LSLS-sequence is exactly LLS-sequence for the corresponding angle. So LSLS-sequence is a natural combinatorial-geometrical generalization of LLS-sequences. Note also that if we know the whole LSLS-sequence for some V -broken line and the coordinates of points $V, A_0,$ and A_1 then the coordinates of A_2, \dots can be restored in the unique way.

Let us show how to identify geometrically the signs of elements of the LSLS-sequence for a lattice oriented V -broken line on Figure 4.




FIGURE 4. All possible (non-degenerate) \mathcal{L}_+ -affine decompositions for angles and segments of a LSLS-sequence.

On Figure 5 we show an example of lattice oriented V -broken line and the corresponding LSLS-sequence.




FIGURE 5. A lattice oriented V -broken line and the corresponding LSLS-sequence.

PROPOSITION 3.3. *A LSLS-sequence for the given lattice oriented broken line and the lattice point is invariant under the group action of the \mathcal{L}_+ -affine transformations.*

3.1.2. On \mathcal{L}_+ -congruence of lattice oriented V -broken lines. Let us formulate necessary and sufficient conditions for two lattice oriented V -broken lines (for the same lattice point V) to be \mathcal{L}_+ -congruent.

THEOREM 3.4. *The LSLS-sequences of two lattice oriented V_1 -broken and V_2 -broken lines (for two lattice points V_1 and V_2) coincide iff there exist a \mathcal{L}_+ -affine transformation taking the point V_1 to V_2 and one lattice oriented broken line to the other.*

PROOF. The LSLS-sequence for any lattice oriented V -broken line is uniquely defined, and by Proposition 3.3 is invariant under the group action of \mathcal{L}_+ -affine orientation preserving transformations. Therefore, the LSLS-sequences for two \mathcal{L}_+ -congruent lattice oriented broken lines coincide.

Suppose now that two lattice oriented V_1 -broken and V_2 -broken lines $A_0 \dots A_n$, and $B_0 \dots B_n$ respectively have the same LSLS-sequence $(a_0, a_1, \dots, a_{2n-3}, a_{2n-2})$. Let us prove that these broken lines are \mathcal{L}_+ -congruent. Without loose of generality we consider the point V_1 at the origin O .

Let ξ be the \mathcal{L}_+ -affine transformation taking the point V_2 to the point $V_1 = O$, B_0 to A_0 , and the lattice straight line containing B_0B_1 to the lattice straight line containing A_0A_1 . Let us prove inductively that $\xi(B_i) = A_i$.

Base of induction. Since $a_0 = b_0$, we have

$$\text{sgn}(A_0OA_1) \ell(A_0A_1) = \text{sgn}(\xi(B_0)O\xi(B_1)) \ell(\xi(B_0)\xi(B_1)).$$

Thus, the lattice segments A_0A_1 and $A_0\xi(B_1)$ are of the same lattice length and of the same direction. Therefore, $\xi(B_1) = A_1$.

Step of induction. Suppose, that $\xi(B_i) = A_i$ holds for any nonnegative integer $i \leq k$, where $k \geq 1$. Let us prove, that $\xi(B_{k+1}) = A_{k+1}$. Denote by C_{k+1} the lattice point $\xi(B_{k+1})$. Let $A_k = (q_k, p_k)$. Denote by A'_k the closest lattice point of the segment $A_{k-1}A_k$ to the vertex A_k . Suppose that $A'_k = (q'_k, p'_k)$. We know also

$$\begin{aligned} a_{2k-1} &= \text{sgn}(A_{k-1}OA_k) \text{sgn}(A_kOC_{k+1}) \text{sgn}(A_{k-1}A_kC_{k+1}) \text{lsin} \angle A_{k-1}A_kC_{k+1}, \\ a_{2k} &= \text{sgn}(A_kOC_{k+1}) \ell(A_kC_{k+1}). \end{aligned}$$

Let the coordinates of C_{k+1} be (x, y) . Since $\ell(A_kC_{k+1}) = |a_{2k}|$ and the segment A_kC_{k+1} is at unit distance to the origin O , we have $\text{IS}(\triangle OA_kC_{k+1}) = |a_{2k}|$. Since the segment OA_k is of the unit lattice length, the coordinates of C_{k+1} satisfy the following equation:

$$|-p_kx + q_ky| = |a_{2k}|.$$

Since $\text{sgn}(A_kOC_{k+1}) \ell(A_kC_{k+1}) = \text{sign}(a_{2k})$, we have $-p_kx + q_ky = a_{2k}$.

Since $\text{lsin} \angle A'_kA_kC_{k+1} = \text{lsin} \angle A_{k-1}A_kC_{k+1} = |a_{2k-1}|$, and the lattice lengths of A_kC_{k+1} , and A'_kA_k are $|a_{2k}|$ and 1 respectively, we have $\text{IS}(\triangle A'_kA_kC_{k+1}) = |a_{2k-1}a_{2k}|$. Therefore, the coordinates of C_{k+1} satisfy the following equation:

$$|-(p_k - p'_k)(x - q_k) + (q_k - q'_k)(y - p_k)| = |a_{2k-1}a_{2k}|.$$

Since

$$\begin{cases} \operatorname{sgn}(A_{k-1}OA_k) \operatorname{sgn}(A_kOC_{k+1}) \operatorname{sgn}(A_{k-1}A_kC_{k+1}) = \operatorname{sign}(a_{2k-1}) \\ \operatorname{sgn}(A_kOC_{k+1}) = \operatorname{sign}(a_{2k}) \end{cases},$$

we have $(p_k - p'_k)(x - q_k) - (q_k - q'_k)(y - p_k) = \operatorname{sgn}(A_{k-1}OA_k)a_{2k-1}a_{2k}$.

We obtain the following:

$$\begin{cases} -p_kx + q_ky = a_{2k} \\ (p_k - p'_k)(x - q_k) - (q_k - q'_k)(y - p_k) = \operatorname{sgn}(A_{k-1}OA_k)a_{2k-1}a_{2k} \end{cases}.$$

Since

$$\left| \det \begin{pmatrix} -p_k & q_k \\ p'_k - p_k & q_k - q'_k \end{pmatrix} \right| = 1,$$

there exist a unique integer solution for the system of equations for x and y . Hence, the points A_{k+1} and C_{k+1} have the same coordinates. Therefore, $\xi(B_{k+1}) = A_{k+1}$. We have proven the step of induction.

The proof of Theorem 3.4 is completed by induction.

3.1.3. Values of continued fractions for lattice oriented broken lines at unit distance from the origin. Now we show the relation between lattice oriented broken lines at unit distance from the origin O and the corresponding continued fractions for them.

THEOREM 3.5. *Let $A_0A_1 \dots A_n$ be a lattice oriented O -broken line. Let also $A_0 = (1, 0)$, $A_1 = (1, a_0)$, $A_n = (p, q)$, where $\gcd(p, q) = 1$, and $(a_0, a_1, \dots, a_{2n-2})$ be the corresponding LSLS-sequence. Then the following holds:*

$$\frac{q}{p} =]a_0, a_1, \dots, a_{2n-2}[.$$

PROOF. To prove this theorem we use an induction on the number of edges of the broken lines.

Base of induction. Suppose that a lattice oriented O -broken line has a unique edge, and the corresponding sequence is (a_0) . Then $A_1 = (1, a_0)$ by the assumptions of the theorem. Therefore, we have $\frac{a_0}{1} =]a_0[$.

Step of induction. Suppose that the statement of the theorem is correct for any lattice oriented O -broken line with k edges. Let us prove the theorem for the arbitrary lattice oriented O -broken line with $k+1$ edges (and satisfying the conditions of the theorem).

Let $A_0 \dots A_{k+1}$ be a lattice oriented O -broken line with the following LSLS-sequence $(a_0, a_1, \dots, a_{2k-1}, a_{2k})$. Let also

$$A_0 = (1, 0), \quad A_1 = (1, a_0), \quad \text{and} \quad A_{k+1} = (p, q).$$

Consider the lattice oriented O -broken line $B_1 \dots B_{k+1}$ with shorter LSLS-sequence for it: $(a_2, a_3, \dots, a_{2k-2}, a_{2k})$. Let also

$$B_1 = (1, 0), \quad B_2 = (1, a_2), \quad \text{and} \quad B_{k+1} = (p', q').$$

By the induction assumption we have

$$\frac{q'}{p'} =]a_2, a_3, \dots, a_{2k}[:$$

We extend the lattice oriented broken line $B_1 \dots B_{k+1}$ to the lattice oriented O -broken line $B_0 B_1 \dots B_{k+1}$, where $B_0 = (1+a_0 a_1, -a_0)$. Let the lattice LSLS-sequence for this broken line be $(b_0, b_1, \dots, b_{2k-1}, b_{2k})$. Note that

$$\begin{aligned} b_0 &= \text{sgn}(B_0 O B_1) |l(B_0 B_1)| = \text{sign}(a_0) |a_0| = a_0, \\ b_1 &= \text{sgn}(B_0 O B_1) \text{sgn}(B_1 O B_2) \text{sgn}(B_0 B_1 B_2) \text{lsin} \angle B_0 B_1 B_2 \\ &= \text{sign } a_0 \text{sign } b_2 \text{sign}(a_0 a_1 b_2) |a_1| = a_1, \\ b_l &= a_l, \quad \text{for } l = 2, \dots, 2k. \end{aligned}$$

Consider a \mathcal{L}_+ -linear transformation ξ that takes the point B_0 to the point $(1, 0)$, and B_1 to $(1, a_0)$. These two conditions uniquely define ξ :

$$\xi = \begin{pmatrix} 1 & a_1 \\ a_0 & 1 + a_0 a_1 \end{pmatrix}.$$

Since $B_{k+1} = (p', q')$, we have $\xi(B_{k+1}) = (p' + a_1 q', q' a_0 + p' + p' a_0 a_1)$.

$$\frac{q' a_1 + p' + p' a_0 a_1}{p' + a_1 q'} = a_0 + \frac{1}{a_1 + q'/p'} =]a_0, a_1, a_2, a_3, \dots, a_{2n}[:$$

Since, by Theorem 3.4 the lattice oriented broken lines $B_0 B_1 \dots B_{k+1}$ and $A_0 A_1 \dots A_{k+1}$ are \mathcal{L} -linear equivalent, $B_0 = A_0$, and $B_1 = A_1$, these broken lines coincide. Therefore, for the coordinates (p, q) the following hold

$$\frac{q}{p} = \frac{q' a_0 + p' + p' a_0 a_1}{p' + a_1 q'} =]a_0, a_1, a_2, a_3, \dots, a_{2k}[:$$

On Figure 6 we illustrate the step of induction with an example of lattice oriented O -broken line with the LSLS-sequence: $(1, -1, 2, 2, -1)$. We start (the left picture) with the broken line $B_1 B_2 B_3$ with the LSLS-sequence: $(2, 2, -1)$. Note that the ratio of the coordinates of the point B_3 is $-3/-1 =]2 : 2; -1[:$. Then, (the picture in the middle) we extend the broken line $B_1 B_2 B_3$ to the broken line $B_0 B_1 B_2 B_3$ with the LSLS-sequence: $(1, -1, 2, 2, -1)$. Finally

(the right picture) we apply a corresponding \mathcal{L}_+ -linear transformation ξ to achieve the resulting broken line $A_0A_1A_2A_3$. Now the ratio of the coordinates of the point A_3 is $-1/2 =]1 : -1; 2; 2; -1[$.




FIGURE 6. The case of lattice oriented O -broken line with LSLS-sequence: $(1, -1, 2, 2, -1)$.

We have proven the step of induction.

The proof of Theorem 3.5 is completed.

REMARK 3.6. Theorem 3.5 immediately implies the statement of Theorem 1.10. One should put the sail of an angle as an oriented-broken line $A_0A_1 \dots A_n$.

3.2. Extended lattice angles. Sums for ordinary and extended lattice angles

3.2.1. Equivalence classes of lattice oriented broken lines and the corresponding extended angles.

DEFINITION 3.7. Consider a lattice point V . Two lattice oriented V -broken lines l_1 and l_2 are said to be *equivalent* if they have in common the first and the last vertices and the closed broken line generated by l_1 and the inverse of l_2 is homotopy equivalent to the point in $\mathbb{R}^2 \setminus \{V\}$.

An equivalence class of lattice oriented V -broken lines containing the broken line $A_0A_1 \dots A_n$ is called the *extended lattice angle for the equivalence class of $A_0A_1 \dots A_n$ at the vertex V* (or, for short, *extended angle*) and denoted by $\angle(V, A_0A_1 \dots A_n)$.

We study the extended angles up to \mathcal{L}_+ -congruence.

DEFINITION 3.8. Two extended angles Φ_1 and Φ_2 are said to be \mathcal{L}_+ -congruent iff there exist a \mathcal{L}_+ -affine transformation sending the class of lattice

oriented broken lines corresponding to Φ_1 to the class of lattice oriented broken lines corresponding to Φ_2 . We denote this by $\Phi_1 \hat{=} \Phi_2$.

3.2.2. *Revolution numbers for extended angles.* Let $r = \{V + \lambda \bar{v} \mid \lambda \geq 0\}$ be the oriented ray for an arbitrary vector \bar{v} with the vertex at V , and AB be an oriented (from A to B) segment not contained in the ray r . Suppose also, that the vertex V of the ray r is not contained in the segment AB . We denote by $\#(r, V, AB)$ the following number:

$$\#(r, V, AB) = \begin{cases} 0, & AB \cap r = \emptyset \\ \frac{1}{2} \operatorname{sgn}(A(A-\bar{v})B), & AB \cap r \in \{A, B\} \\ \operatorname{sgn}(A(A-\bar{v})B), & AB \cap r \in AB \setminus \{A, B\}, \end{cases}$$

and call it the *intersection number* of the ray r and the segment AB .

DEFINITION 3.9. Let $A_0A_1 \dots A_n$ be some lattice oriented broken line, and let r be an oriented ray $\{V + \lambda \bar{v} \mid \lambda \geq 0\}$. Suppose that the ray r does not contain the edges of the broken line, and the broken line does not contain the point V . We call the number

$$\sum_{i=1}^n \#(r, V, A_{i-1}A_i)$$

the *intersection number* of the ray r and the lattice oriented broken line $A_0A_1 \dots A_n$, and denote it by $\#(r, V, A_0A_1 \dots A_n)$.

DEFINITION 3.10. Consider an arbitrary extended angle $\angle(V, A_0A_1 \dots A_n)$. Denote the rays $\{V + \lambda \overline{VA_0} \mid \lambda \geq 0\}$ and $\{V - \lambda \overline{VA_0} \mid \lambda \geq 0\}$ by r_+ and r_- respectively. The number

$$\frac{1}{2}(\#(r_+, V, A_0A_1 \dots A_n) + \#(r_-, V, A_0A_1 \dots A_n))$$

is called the *lattice revolution number* for the extended angle $\angle(V, A_0A_1 \dots A_n)$, and denoted by $\#(\angle(V, A_0A_1 \dots A_n))$. We say also that $\#(\angle(V, A_0)) = 0$.

Let us give some examples. Let $O = (0, 0)$, $A = (1, 0)$, $B = (0, 1)$, $C = (-1, -1)$, then

$$\begin{aligned} \#(\angle(O, A)) &= 0, & \#(\angle(O, AB)) &= \frac{1}{4}, \\ \#(\angle(O, ABCA)) &= 1, & \#(\angle(O, ACB)) &= -\frac{3}{4}. \end{aligned}$$

Now we show that the definition of revolution number is correct.

PROPOSITION 3.11. *The revolution number of any extended angle is well-defined.*

PROOF. Consider an arbitrary extended angle $\angle(V, A_0A_1 \dots A_n)$. Let

$$r_+ = \{V + \lambda \overline{VA_0} \mid \lambda \geq 0\} \quad \text{and} \quad r_- = \{V - \lambda \overline{VA_0} \mid \lambda \geq 0\}.$$

Since the lattice oriented broken line $A_0A_1 \dots A_n$ is at unit distance from the point V , any segment of this broken line is at unit distance from V . Thus, the broken line does not contain V , and the rays r_+ and r_- do not contain edges of the curve.

Suppose that

$$\angle(V, A_0A_1 \dots A_n) = \angle(V', A'_0A'_1 \dots A'_m).$$

This implies that $V = V'$, $A_0 = A'_0$, $A_n = A'_m$, and the broken line $A_0A_1 \dots A_nA'_{m-1} \dots A'_1A'_0$ is homotopy equivalent to the point in $\mathbb{R}^2 \setminus \{V\}$. Thus,

$$\begin{aligned} \#(\angle(V, A_0A_1 \dots A_n)) - \#(\angle(V', A'_0A'_1 \dots A'_m)) \\ &= \frac{1}{2}(\#(r_+, V, A_0A_1 \dots A_nA'_{m-1} \dots A'_1A'_0) \\ &\quad + \#(r_-, V, A_0A_1 \dots A_nA'_{m-1} \dots A'_1A'_0)) \\ &= 0 + 0 = 0. \end{aligned}$$

Hence,

$$\#(\angle(V, A_0A_1 \dots A_n)) = \#(\angle(V', A'_0A'_1 \dots A'_m)).$$

Therefore, the revolution number of any extended angle is well-defined.

PROPOSITION 3.12. *The revolution number of extended angles is invariant under the group action of the \mathcal{L}_+ -affine transformations.*

3.2.3. *Zero ordinary angles.* For the next theorem we will need to define zero ordinary angles and their trigonometric functions. Let A , B , and C be three lattice points of the same lattice straight line. Suppose that B is distinct to A and C and the rays BA and BC coincide. We say that the ordinary angle with the vertex at B and the rays BA and BC is *zero*. Suppose $\angle ABC$ is zero, put by definition

$$\text{l sin}(\angle ABC) = 0, \quad \text{l cos}(\angle ABC) = 1, \quad \text{l tan}(\angle ABC) = 0.$$

Denote by $\text{larctan}(0)$ the angle $\angle AOA$ where $A = (1, 0)$, and O is the origin.


3.2.4. *On normal forms of extended angles.* Let us formulate and prove a theorem on normal forms of extended angles. We use the following notation: by the sequence

$$((a_0, \dots, a_n) \times k\text{-times}, b_0, \dots, b_m),$$

where $k \geq 0$, we denote the following sequence:


$$\underbrace{(a_0, \dots, a_n, a_0, \dots, a_n, \dots, a_0, \dots, a_n)}_{k\text{-times}}, b_0, \dots, b_m).$$

DEFINITION 3.13. I) Suppose O be the origin, A_0 be the point $(1, 0)$. We say that the extended angle $\angle(O, A_0)$ is of the type I and denote it by $0\pi + \text{larctan}(0)$ (or 0 , for short). The empty sequence is said to be *characteristic* for the angle $0\pi + \text{larctan}(0)$.




Consider a lattice oriented O -broken line $A_0A_1 \dots A_s$, where O is the origin. Let also A_0 be the point $(1, 0)$, and the point A_1 be on the straight line $x = 1$. If the LSLS-sequence of the extended angle $\Phi_0 = \angle(O, A_0A_1 \dots A_s)$ coincides with the following sequence (we call it *characteristic sequence* for the corresponding angle):


$\text{II}_k) ((1, -2, 1, -2) \times (k - 1)\text{-times}, 1, -2, 1)$, where $k \geq 1$, then we denote the angle Φ_0 by $k\pi + \text{larctan}(0)$ (or $k\pi$, for short) and say that Φ_0 is of the type II_k ;




$\text{III}_k) ((-1, 2, -1, 2) \times (k - 1)\text{-times}, -1, 2, -1)$, where $k \geq 1$, then we denote the angle Φ_0 by $-k\pi + \text{larctan}(0)$ (or $-k\pi$, for short) and say that Φ_0 is of the type III_k ;



IV_k) $((1, -2, 1, -2) \times k\text{-times}, a_0, \dots, a_{2n})$, where $k \geq 0, n \geq 0, a_i > 0$, for $i = 0, \dots, 2n$, then we denote the angle Φ_0 by $k\pi + \text{larctan}([a_0, a_1, \dots, a_{2n}])$ and say that Φ_0 is of the type IV_k;



V_k) $((-1, 2, -1, 2) \times k\text{-times}, a_0, \dots, a_{2n})$, where $k > 0, n \geq 0, a_i > 0$, for $i = 0, \dots, 2n$, then we denote the angle Φ_0 by $-k\pi + \text{larctan}([a_0, a_1, \dots, a_{2n}])$ and say that Φ_0 is of the type V_k.



THEOREM 3.14. For any extended angle Φ there exist a unique type among the types I–V and a unique extended angle Φ_0 of that type such that Φ_0 is \mathcal{L}_+ -congruent to Φ .

The extended angle Φ_0 is said to be the normal form for the extended angle Φ .

For the proof of Theorem 3.14 we need the following lemma.

LEMMA 3.15. Let $m, k \geq 1$, and $a_i > 0$ for $i = 0, \dots, 2n$ be some integers.

a) Suppose the LSLS-sequences for the extended angles Φ_1 and Φ_2 are respectively

$$((1, -2, 1, -2) \times (k-1)\text{-times}, 1, -2, 1, -2, a_0, \dots, a_{2n})$$

and

$$((1, -2, 1, -2) \times (k-1)\text{-times}, 1, -2, 1, m, a_0, \dots, a_{2n}),$$

then Φ_1 is \mathcal{L}_+ -congruent to Φ_2 .

b) Suppose the LSLS-sequences for the extended angles Φ_1 and Φ_2 are respectively

$$((-1, 2, -1, 2) \times (k-1)\text{-times}, -1, 2, -1, m, a_0, \dots, a_{2n})$$

and

$$((-1, 2, -1, 2) \times (k-1)\text{-times}, -1, 2, -1, 2, a_0, \dots, a_{2n}),$$

then Φ_1 is \mathcal{L}_+ -congruent to Φ_2 .

PROOF. We prove the first statement of the lemma. Suppose that m is integer, k is positive integer, and a_i for $i = 0, \dots, 2n$ are positive integers.

Let us construct the angle Ψ_1 with vertex at the origin for the lattice oriented broken line $A_0 \dots A_{2k+n+1}$, corresponding to the LSLS-sequence

$$((1, -2, 1, -2) \times (k-1)\text{-times}, 1, -2, 1, -2, a_0, \dots, a_{2n}),$$

such that $A_0 = (1, 0)$, $A_1 = (1, 1)$. Note that

$$\begin{cases} A_{2l} = ((-1)^l, 0), & \text{for } l < k - 1 \\ A_{2l+1} = ((-1)^l, (-1)^l), & \text{for } l < k - 1 \\ A_{2k} = ((-1)^k, 0) \\ A_{2k+1} = ((-1)^k, (-1)^k a_0) \end{cases}.$$

Let us construct the angle Ψ_2 with vertex at the origin for the lattice oriented broken line $B_0 \dots B_{2k+n+1}$, corresponding to the LSLS-sequence

$$((1, -2, 1, -2) \times (k-1)\text{-times}, 1, -2, 1, m, a_0, \dots, a_{2n}).$$

such that $B_0 = (1, 0)$, $B_1 = (-m - 1, 1)$. Note also that

$$\begin{cases} B_{2l} = ((-1)^l, 0), & \text{for } l < k - 1 \\ B_{2l+1} = ((-1)^l(-m - 1), (-1)^l), & \text{for } l < k - 1 \\ B_{2k} = ((-1)^k, 0) \\ B_{2k+1} = ((-1)^k, (-1)^k a_0) \end{cases}.$$

From the above we know, that the points A_{2k} and A_{2k+1} coincide with the points B_{2k} and B_{2k+1} respectively. Since the remaining parts of both LSLS-sequences (i. e. (a_0, \dots, a_{2n})) coincide, the point A_l coincide with the point B_l for $l > 2k$.

Since the lattice oriented broken lines $A_0 \dots A_{2k}$ and $B_0 \dots B_{2k}$ are of the same equivalence class, and the point A_l coincide with the point B_l for $l > 2k$, we obtain

$$\Psi_1 = \angle(O, A_0 \dots A_{2k+n+1}) = \angle(O, B_0 \dots B_{2k+n+1}) = \Psi_2.$$

Therefore, by Theorem 3.4 we have the following:

$$\Phi_1 \hat{\cong} \Psi_1 = \Psi_2 \hat{\cong} \Phi_2.$$

This concludes the proof of Lemma 3.15a.

Since the proof of Lemma 3.15b almost completely repeats the proof of Lemma 3.15a, we omit the proof of Lemma 3.15b here.

PROOF OF THEOREM 3.14. First, we prove that any two distinct extended angles listed in Definition 3.13 are not \mathcal{L}_+ -congruent. Let us note that the revolution numbers of extended angles distinguish the types of the angles. The revolution number for the extended angle of the type I is 0. The revolution number for the extended angle of the type II_k is $1/2(k + 1)$ where $k \geq 0$. The revolution number for the extended angle of the type III_k is $-1/2(k + 1)$ where $k \geq 0$. The revolution number for the extended angles of the type IV_k is $1/4 + 1/2k$ where $k \geq 0$. The revolution number for the extended angles of the type V_k is $1/4 - 1/2k$ where $k > 0$.

So we have proven that two extended angles of different types are not \mathcal{L}_+ -congruent. For the types I, II_k , and III_k the proof is completed, since any such type consists of the unique extended angle.

Let us prove that normal forms of the same type IV_k (or of the same type V_k) are not \mathcal{L}_+ -congruent for any integer $k \geq 0$ (or $k > 0$). Consider an extended angle $\Phi = k\pi + \text{larctan}([a_0, a_1, \dots, a_{2n}])$. Suppose that a lattice oriented O -broken line $A_0A_1 \dots A_m$, where $m = 2|k| + n + 1$ defines the angle Φ . Let also that the LSLS-sequence for this broken line be characteristic.

Suppose, that k is even, then the ordinary angle $\angle A_0OA_m$ is \mathcal{L}_+ -congruent to the ordinary angle $\text{larctan}([a_0, a_1, \dots, a_{2n}])$. This angle is a \mathcal{L}_+ -affine invariant for the extended angle Φ . This invariant distinguish the extended angles of type IV_k (or V_k) with even k .

Suppose, that k is odd, then denote $B = O + \overline{A_0O}$. The ordinary angle $\angle BVA_m$ is \mathcal{L}_+ -congruent to the ordinary angle $\text{larctan}([a_0, a_1, \dots, a_{2n}])$. This angle is a \mathcal{L}_+ -affine invariant for the extended angle Φ . This invariant distinguish the extended angles of type IV_k (or V_k) with odd k .

Therefore, the extended angles listed in Definition 3.13 are not \mathcal{L}_+ -congruent.

Now we prove that an arbitrary extended angle is \mathcal{L}_+ -congruent to one of the extended angles of the types I–V.

Consider an arbitrary extended angle $\angle(V, A_0A_1 \dots A_n)$ and denote it by Φ . If $\#(\Phi) = k/2$ for some integer k , then Φ is \mathcal{L}_+ -congruent to an angle of one of the types I–III. Let $\#(\Phi) = 1/4$, then the extended angle Φ is \mathcal{L}_+ -congruent to the extended angle defined by the sail of the ordinary angle $\angle A_0VA_n$ of the type IV_0 .

Suppose now, that $\#(\Phi) = 1/4 + k/2$ for some positive integer k , then one of its LSLS-sequence is of the following form:

$$((1, -2, 1, -2) \times (k - 1)\text{-times}, 1, -2, 1, m, a_0, \dots, a_{2n}),$$

where $a_i > 0$, for $i = 0, \dots, 2n$. By Lemma 3.15 the extended angle defined by this sequence is \mathcal{L}_+ -congruent to an extended angle of the type IV_k defined by the sequence

$$((1, -2, 1, -2) \times (k - 1)\text{-times}, 1, -2, 1, -2, a_0, \dots, a_{2n}).$$

Finally, let $\#(\Phi) = 1/4 - k/2$ for some positive integer k , then one of its LSLS-sequence is of the following form:

$$((-1, 2, -1, 2) \times (k - 1)\text{-times}, -1, 2, -1, m, a_0, \dots, a_{2n}),$$

where $a_i > 0$, for $i = 0, \dots, 2n$. By Lemma 3.15 the extended angle defined by this sequence is \mathcal{L}_+ -congruent to an extended angle of the type V_k defined by the sequence

$$((-1, 2, -1, 2) \times (k - 1)\text{-times}, -1, 2, -1, 2, a_0, \dots, a_{2n}).$$

This completes the proof of Theorem 3.14.

Let us finally give the definition of trigonometric functions for the extended angles and describe some relations between ordinary and extended angles.

DEFINITION 3.16. Consider an arbitrary extended angle Φ with the normal form $k\pi + \varphi$ for some ordinary (possible zero) angle φ and for an integer k .

- a) The ordinary angle φ is said to be *associated* with the extended angle Φ .
- b) The numbers $\text{ltan}(\varphi)$, $\text{lsin}(\varphi)$, and $\text{lcos}(\varphi)$ are called the *lattice tangent*, the *lattice sine*, and the *lattice cosine* of the extended angle Φ .

Since all sails for ordinary angles are lattice oriented broken lines, the set of all ordinary angles is naturally embedded into the set of extended angles.

DEFINITION 3.17. For an ordinary angle φ the angle

$$0\pi + \text{larctan}(\text{ltan } \varphi)$$

is said to be *corresponding* to the angle φ and denoted by $\bar{\varphi}$.

From Theorem 3.14 it follows that for every ordinary angle φ there exists and unique an extended angle $\bar{\varphi}$ corresponding to φ . Therefore, two ordinary angles φ_1 and φ_2 are \mathcal{L} -congruent iff the corresponding lattice angles $\bar{\varphi}_1$ and $\bar{\varphi}_2$ are \mathcal{L}_+ -congruent.

3.2.5. *Opposite extended angles. Sums of extended angles. Sums of ordinary angles.* Consider an extended angle Φ with the vertex V for some equivalence class of a given lattice oriented broken line. The extended angle Ψ with the vertex V for the equivalence class of the inverse lattice oriented broken line is called *opposite* to the given one and denoted by $-\Phi$.

PROPOSITION 3.18. *For any extended angle $\Phi \hat{=} k\pi + \varphi$ we have:*

$$-\Phi \hat{=} (-k - 1)\pi + (\pi - \varphi).$$

Let us introduce the definition of sums of ordinary and extended angles.

DEFINITION 3.19. Consider arbitrary extended angles $\Phi_i, i = 1, \dots, l$. Let the characteristic sequences for the normal forms of Φ_i be $(a_{0,i}, a_{1,i}, \dots, a_{2n_i,i})$ for $i = 1, \dots, l$. Let $M = (m_1, \dots, m_{l-1})$ be some $(l - 1)$ -tuple of integers. The normal form of any extended angle, corresponding to the following LSLS-sequence

$$(a_{0,1}, a_{1,1}, \dots, a_{2n_1,1}, m_1, a_{0,2}, \dots, a_{2n_2,2}, m_2, \dots, m_{l-1}, a_{0,l}, \dots, a_{2n_l,l}),$$

is called the *M-sum of extended angles Φ_i ($i = 1, \dots, l$)* and denoted by

$$\sum_{M,i=1}^l \Phi_i, \quad \text{or equivalently by} \quad \Phi_1 +_{m_1} \Phi_2 +_{m_2} \dots +_{m_{l-1}} \Phi_l.$$

PROPOSITION 3.20. *The M-sum of extended angles Φ_i ($i = 1, \dots, l$) is well-defined.*

Let us say a few words about properties of *M*-sums.

Notice that *M*-sum of extended angles is non-associative. For example, let $\Phi_1 \hat{=} \text{larctan } 2$, $\Phi_2 \hat{=} \text{larctan}(3/2)$, and $\Phi_3 \hat{=} \text{larctan } 5$. Then

$$\begin{aligned} \Phi_1 +_{-1} \Phi_2 +_{-1} \Phi_3 &= \pi + \text{larctan}(4), \\ \Phi_1 +_{-1} (\Phi_2 +_{-1} \Phi_3) &= 2\pi, \\ (\Phi_1 +_{-1} \Phi_2) +_{-1} \Phi_3 &= \text{larctan}(1). \end{aligned}$$

The M -sum of extended angles is non-commutative. For example, let $\Phi_1 \hat{\cong} \text{larctan } 1$, and $\Phi_2 \hat{\cong} \text{larctan } 5/2$. Then

$$\Phi_1 +_1 \Phi_2 = \text{larctan}(12/7) \neq \text{larctan}(13/5) = \Phi_2 +_1 \Phi_1.$$

REMARK 3.21. The M -sum of extended angles is naturally extended to the sum of classes of \mathcal{L}_+ -congruences of extended angles.

We conclude this section with the definition of sums of ordinary angles.

DEFINITION 3.22. Consider ordinary angles α_i , where $i = 1, \dots, l$. Let $\bar{\alpha}_i$ be the corresponding extended angles for α_i , and $M = (m_1, \dots, m_{l-1})$ be some $(l-1)$ -tuple of integers. The ordinary angle φ associated with the extended angle

$$\Phi = \bar{\alpha}_1 +_{m_1} \bar{\alpha}_2 +_{m_2} \cdots +_{m_{l-1}} \bar{\alpha}_l.$$

is called the M -sum of ordinary angles α_i ($i = 1, \dots, l$) and denoted by

$$\sum_{M, i=1}^l \alpha_i, \quad \text{or equivalently by } \alpha_1 +_{m_1} \alpha_2 +_{m_2} \cdots +_{m_{l-1}} \alpha_l.$$

REMARK 3.23. Note that the sum of ordinary angles is naturally extended to the classes of \mathcal{L} -congruences of lattice angles.

4. Relations between extended and ordinary lattice angles. Proof of the first statement of Theorem 2.2

Throughout this section we again fix some lattice basis and use the system of coordinates OXY corresponding to this basis.

4.1. On relations between continued fractions for lattice oriented broken lines and the lattice tangents of the corresponding extended angles

For a real number r we denote by $\lfloor r \rfloor$ the maximal integer not greater than r .

THEOREM 4.1. Consider an extended angle $\Phi = \angle(V, A_0A_1 \dots A_n)$. Suppose, that the normal form for Φ is $k\pi + \varphi$ for some integer k and an ordinary angle φ . Let $(a_0, a_1, \dots, a_{2n-2})$ be the LSLS-sequence for the lattice oriented broken line $A_0A_1 \dots A_n$. Suppose that

$$\lfloor a_0, a_1, \dots, a_{2n-2} \rfloor = q/p.$$

Then the following holds:

$$\varphi \cong \begin{cases} \text{larctan}(1), & \text{if } q/p = \infty \\ \text{larctan}(q/p), & \text{if } q/p \geq 1 \\ \text{larctan}\left(\frac{|q|}{|p| - \lfloor (|p|-1)/|q| \rfloor |q|}\right), & \text{if } 0 < q/p < 1 \\ 0, & \text{if } q/p = 0 \\ \pi - \text{larctan}\left(\frac{|q|}{|p| - \lfloor (|p|-1)/|q| \rfloor |q|}\right), & \text{if } -1 < q/p < 0 \\ \pi - \text{larctan}(-q/p), & \text{if } q/p \leq -1 \end{cases}.$$

PROOF. Consider the following linear coordinates $(*, *)'$ on the plane, associated with the lattice oriented V -broken line $A_0A_1 \dots A_n$. Let the origin O' be at the vertex V , $(1, 0)' = A_0$, and $(1, 1)' = A_0 + \frac{1}{a_0} \text{sgn}(A_0O'A_1)A_0A_1$. The other coordinates are uniquely defined by linearity. We denote this system of coordinates by $O'X'Y'$.

The set of integer points for the coordinate system $O'X'Y'$ coincides with the set of lattice points of the plane. The basis of vectors $(1, 0)'$ and $(0, 1)'$ defines a positive orientation.

Suppose that the new coordinates of the point A_n are $(p', q)'$. Then by Theorem 3.5 we have $q'/p' = q/p$. This directly implies the statement of the theorem for the cases $q' > p' > 0$, $q'/p' = 0$, and $q'/p' = \infty$.

Suppose now that $p' > q' > 0$. Consider the ordinary angle $\varphi = \angle A_0PA_n$. Let $B_0 \dots B_m$ be the sail for it. The direct calculations show that the point

$$D = B_0 + \frac{\overline{B_0B_1}}{\ell(B_0B_1)}$$

coincides with the point $(1 + \lfloor (p' - 1)/q' \rfloor, 1)$ in the system of coordinates $O'X'Y'$.

Consider the \mathcal{L}_+ -linear (in the coordinates $O'X'Y'$) transformation ξ that takes the point $A_0 = B_0$ to itself, and the point D to $(1, 1)'$. These conditions uniquely identify ξ .

$$\xi = \begin{pmatrix} 1 & -\lfloor (p' - 1)/q' \rfloor \\ 0 & 1 \end{pmatrix}$$

The transformation ξ takes the point $A_n = B_m$ with the coordinates $(p', q)'$ to the point with the coordinates $(p' - \lfloor (p' - 1)/q' \rfloor, q)'$. Since $q'/p' = q/p$, we obtain the following

$$\varphi = \text{larctan}\left(\frac{q'}{p' - \lfloor (p' - 1)/q' \rfloor q'}\right) = \text{larctan}\left(\frac{q}{p - \lfloor (p - 1)/q \rfloor q}\right).$$

The proof for the case $q' > 0$ and $p' < 0$ repeats the described cases after taking to the consideration the adjacent angles.

Finally, the case of $q' < 0$ repeats all previous cases by the central symmetry (centered at the point O') reasons.

This completes the proof of Theorem 4.1.

COROLLARY 4.2. *The revolution number and the continued fraction for a lattice oriented broken line at unit distance from the vertex uniquely define the \mathcal{L}_+ -congruence class of the corresponding extended angle.*

4.2. Proof of Theorem 2.2a: two preliminary lemmas

We say that the lattice point P is at lattice distance k from the lattice segment AB if the lattice vectors of the segment AB and the vector \overline{AP} generate a sublattice of the lattice of index k .

DEFINITION 4.3. Consider a lattice triangle $\triangle ABC$. Denote the number of lattice points at unit lattice distance from the segment AB and contained in the (closed) triangle $\triangle ABC$ by $l_1(AB; C)$ (see on Figure 7).

Note that all lattice points at lattice unit distance from the segment AB in the (closed) lattice triangle $\triangle ABC$ are contained in one straight line parallel to the straight line AB . Besides, the integer $l_1(AB; C)$ is positive for any triangle $\triangle ABC$.




FIGURE 7. For the given triangle $\triangle ABC$ we have $l_1(AB; C) = 5$.

Now we prove the following lemma.

LEMMA 4.4. *For any lattice triangle $\triangle ABC$ the following holds*

$$\overline{\angle CAB} + l_1(AB) - l_1(AB; C) - 1 \overline{\angle ABC} + l_1(BC) - l_1(BC; A) - 1 \overline{\angle BCA} = \pi.$$

PROOF. Consider an arbitrary lattice triangle $\triangle ABC$. Suppose that the pair of vectors \overline{BA} and \overline{BC} defines the positive orientation of the plane (otherwise we apply to the triangle $\triangle ABC$ some \mathcal{L} -affine transformation changing the

orientation and come to the same position). Denote (see Figure 9 below): $D = A + \overline{BC}$, and $E = A + \overline{AC}$.

Since $CADB$ is a parallelogram, the triangle $\triangle BAD$ is \mathcal{L}_+ -congruent to the triangle $\triangle ABC$. Thus, the angle $\angle BAD$ is \mathcal{L}_+ -congruent to the angle $\angle ABC$, and $l\ell_1(BA; D) = l\ell_1(AB; C)$. Since $EABD$ is a parallelogram, the triangle $\triangle AED$ is \mathcal{L}_+ -congruent to the triangle $\triangle BAD$, and hence is \mathcal{L}_+ -congruent to the triangle $\triangle ABC$. Thus, $\angle DAE$ is \mathcal{L}_+ -congruent to $\angle BCA$, and $l\ell_1(DA; E) = l\ell_1(BC; A)$.

Let $A_0 \dots A_n$ be the sail of $\angle CAB$ with the corresponding LLS-sequence (a_0, \dots, a_{2n-2}) . Let $B_0 B_1 \dots B_m$ be the sail of $\angle BAD$ (where $B_0 = A_n$) with the corresponding LLS-sequence (b_0, \dots, b_{2m-2}) . And let $C_0 C_1 \dots C_l$ be the sail of $\angle DAE$ (where $C_0 = B_m$) with the corresponding LLS-sequence (c_0, \dots, c_{2l-2}) .

Consider now the lattice oriented broken line

$$A_0 \dots A_n B_1 B_2 \dots B_m C_1 C_2 \dots C_l.$$

The LSLS-sequence for this broken line is

$$(a_0, \dots, a_{2n-2}, t, b_0, \dots, b_{2m-2}, u, c_0, \dots, c_{2l-2}),$$

where integers t and u are integers defined by the broken line. By definition of the sum of extended angles this sequence defines the extended angle

$$\overline{\angle CAB} +_t \overline{\angle BAD} +_u \overline{\angle DAE}.$$

By Theorem 4.1, we have $\overline{\angle CAB} +_t \overline{\angle BAD} +_u \overline{\angle DAE} = \pi$.

We compute now the integer t . Denote by A'_n the closest lattice point to the point A_n and distinct to A_n in the segment $A_{n-1}A_n$. Consider the set of lattice points at unit lattice distance from the segment AB and lying in the half-plane with the boundary straight line AB and containing the point D . This set coincides with the following set (See Figure 8):

$$\{A_{n,k} = A_n + \overline{A'_n A_n} + k \overline{A A_n} \mid k \in \mathbf{Z}\}.$$

Since $A_{n,-2} = A + \overline{A'_n A}$, the points $A_{n,k}$ for $k \leq -2$ are in the closed half-plane bounded by the straight line AC and not containing the point B .

Since $A_{n,-1} = A + \overline{A'_n A_n}$, the points $A_{n,k}$ for $k \geq -1$ are in the open half-plane bounded by the straight line AC and containing the point B .

The intersection of the parallelogram $AEDB$ and the open half-plane bounded by the straight line AC and containing the point B contains exactly $l\ell(AB)$ points of the described set: only the points $A_{n,k}$ with $-1 \leq k \leq l\ell(AB) - 2$.




FIGURE 8. Lattice points $A_{n,t}$.

Since the triangle $\triangle BAD$ is \mathcal{L}_+ -congruent to $\triangle ABC$, the number of points $A_{n,k}$ in the closed triangle $\triangle BAD$ is $l\ell_1(AB; C)$: the points $A_{n,k}$ for

$$l\ell(AB) - l\ell_1(AB; C) - 1 \leq k \leq l\ell(AB) - 2.$$

Denote the integer $l\ell(AB) - l\ell_1(AB; C) - 1$ by k_0 .

The point A_{n,k_0} is contained in the segment B_0B_1 of the sail for the ordinary angle $\angle BAD$ (see Figure 9). Since the angles $\angle BAD$ and $\angle ABC$ are \mathcal{L}_+ -congruent, we have

$$\begin{aligned} t &= \operatorname{sgn}(A_{n-1}A_n) \operatorname{sgn}(A_nAB_1) \operatorname{sgn}(A_{n-1}A_nB_1) \operatorname{l}\sin \angle A_{n-1}A_nB_1 \\ &= 1 \cdot 1 \cdot \operatorname{sgn}(A_{n-1}A_nA_{n,k_0}) \operatorname{l}\sin \angle A_{n-1}A_nA_{n,k_0} \\ &= \operatorname{sign}(k_0)|k_0| = k_0 = l\ell(AB) - l\ell_1(AB; C) - 1. \end{aligned}$$




FIGURE 9. The point A_{n,k_0} .

Exactly by the same reasons,

$$u = l\ell(DA) - l\ell_1(DA; E) - 1 = l\ell(BC) - l\ell_1(BC; A) - 1.$$

Therefore, $\overline{\angle CAB} +_{l\ell(AB)-l\ell_1(AB;C)-1} \overline{\angle ABC} +_{l\ell(BC)-l\ell_1(BC;A)-1} \overline{\angle BCA} = \pi$.

LEMMA 4.5. *Let α , β , and γ be nonzero ordinary angles. Suppose that $\bar{\alpha} +_u \bar{\beta} +_v \bar{\gamma} = \pi$, then there exist a triangle with three consecutive ordinary angles \mathcal{L} -congruent to α , β , and γ .*

PROOF. Denote by O the point $(0, 0)$, by A the point $(1, 0)$, and by D the point $(-1, 0)$ in the fixed system of coordinates OXY .

Let us choose the points $B = (p_1, q_1)$ and $C = (p_2, q_2)$ with integers p_1, p_2 and positive integers q_1, q_2 such that

$$\angle AOB = \text{Iarctan}(\text{Itan } \alpha), \quad \text{and} \quad \angle AOC = \overline{\angle AOB} +_u \bar{\beta}.$$

Thus the vectors \overline{OB} and \overline{OC} defines the positive orientation, and $\angle BOC \cong \beta$. Since

$$\bar{\alpha} +_u \bar{\beta} +_v \bar{\gamma} = \pi \quad \text{and} \quad \bar{\alpha} +_u \bar{\beta} \hat{\cong} \angle AOC,$$

the ordinary angle $\angle COD$ is \mathcal{L} -congruent to γ .

Denote by B' the point (p_1q_2, q_1q_2) , and by C' the point (p_2q_1, q_1q_2) and consider the triangle $B'OC'$. Since the ordinary angle $\angle B'OC'$ coincides with the ordinary angle $\angle BOC$, we obtain

$$\angle B'OC' \cong \beta.$$

Since the ordinary angle β is nonzero, the points B' and C' are distinct and the straight line $B'C'$ does not coincide with the straight line OA . Since the second coordinate of the both points B' and C' equal q_1q_2 , the straight line $B'C'$ is parallel to the straight line OA . Thus, by Proposition 1.13 it follows that

$$\angle C'B'O \cong \angle AOB' = \angle AOB \cong \alpha, \quad \text{and} \quad \angle OC'B' \cong \angle C'OD = \angle COD \cong \gamma.$$

So, we have constructed the triangle $\triangle B'OC'$ with three consecutive ordinary angles \mathcal{L} -congruent to α , β , and γ .

4.3. Proof of Theorem 2.2a: conclusion of the proof

Now we return to the proof of the first statement of the theorem on sums of lattice tangents for ordinary angles in lattice triangles.

PROOF OF THEOREM 2.2A. Let α , β , and γ be nonzero ordinary angles satisfying the conditions i) and ii) of Theorem 2.2a.

The second condition: $|\text{Itan}(\alpha), -1, \text{Itan}(\beta), -1, \text{Itan}(\gamma)| = 0$ implies that

$$\bar{\alpha} +_{-1} \bar{\beta} +_{-1} \bar{\gamma} = k\pi.$$

Since all three tangents are positive, we have $k = 1$, or $k = 2$.

Consider the first condition: $]\lvert \tan \alpha, -1, \lvert \tan \beta [$ is either negative or greater than $\lvert \tan \alpha$. It implies that $\bar{\alpha} +_{-1} \bar{\beta} = 0\pi + \varphi$, for some ordinary angle φ , and hence $k = 1$.

Therefore, by Lemma 4.5 there exist a triangle with three consecutive ordinary angles \mathcal{L} -congruent to α, β , and γ .

Let us prove the converse. We prove that condition ii) of Theorem 2.2a holds by reductio ad absurdum. Suppose, that there exist a triangle $\triangle ABC$ with consecutive ordinary angles $\alpha = \angle CAB, \beta = \angle ABC$, and $\gamma = \angle BCA$, such that

$$\begin{cases}]\lvert \tan(\alpha), -1, \lvert \tan(\beta), -1, \lvert \tan(\gamma) [\neq 0 \\]\lvert \tan(\beta), -1, \lvert \tan(\gamma), -1, \lvert \tan(\alpha) [\neq 0 . \\]\lvert \tan(\gamma), -1, \lvert \tan(\alpha), -1, \lvert \tan(\beta) [\neq 0 \end{cases}$$

These inequalities and Lemma 4.4 imply that at least two of the integers

$$\begin{aligned} & \lvert \ell(AB) - \lvert \ell_1(AB; C) - 1, \\ & \lvert \ell(BC) - \lvert \ell_1(BC; A) - 1, \end{aligned}$$

and

$$\lvert \ell(CA) - \lvert \ell_1(CA; B) - 1$$

are nonnegative.

Without losses of generality we suppose that

$$\begin{cases} \lvert \ell(AB) - \lvert \ell_1(AB; C) - 1 \geq 0 \\ \lvert \ell(BC) - \lvert \ell_1(BC; A) - 1 \geq 0 . \end{cases}$$

Since all integers of the continued fraction

$$r =]\lvert \tan(\alpha), \lvert \ell(AB) - \lvert \ell_1(AB; C) - 1, \lvert \tan(\beta), \lvert \ell(BC) - \lvert \ell_1(BC; A) - 1, \lvert \tan(\gamma) [$$

are non-negative and the last one is positive, we obtain that $r > 0$ (or $r = \infty$). From the other hand, by Lemma 4.4 and by Theorem 4.1 we have that $r = 0/-1 = 0$. We come to the contradiction.

Now we prove that condition i) of Theorem 2.2a holds. Suppose that there exist a triangle $\triangle ABC$ with consecutive ordinary angles $\alpha = \angle CAB, \beta = \angle ABC$, and $\gamma = \angle BCA$, such that

$$]\lvert \tan(\alpha), -1, \lvert \tan(\beta), -1, \lvert \tan(\gamma) [= 0.$$

Since $\bar{\alpha} +_{-1} \bar{\beta} +_{-1} \bar{\gamma} = \pi$, we have $\bar{\alpha} +_{-1} \bar{\beta} = 0\pi + \varphi$ for some ordinary angle φ . Therefore, the first condition of the theorem holds.

This concludes the proof of Theorem 2.2.

4.4. *Theorem on sum of lattice tangents for ordinary lattice angles of convex polygons*

A satisfactory description for \mathcal{L} -congruence classes of lattice convex polygons has not been yet found. It is only known that the number of convex polygons with lattice area bounded from above by n grows exponentially in $n^{1/3}$, while n tends to infinity (see [2] and [3]). We conclude this section with the following theorem on necessary and sufficient condition for the lattice angles to be the angles of some convex lattice polygon.

THEOREM 4.6. *Let $\alpha_1, \dots, \alpha_n$ be an arbitrary ordered n -tuple of ordinary non-zero (lattice) angles. Then the following two conditions are equivalent:*

- *there exist a convex n -vertex polygon with consecutive ordinary angles \mathcal{L} -congruent to the ordinary angles α_i for $i = 1, \dots, n$;*
- *there exist a set of integers $M = \{m_1, \dots, m_{n-1}\}$ such that*

$$\sum_{M, i=1}^n \overline{\pi - \alpha_i} = 2\pi.$$

PROOF. Consider an arbitrary n -tuple of ordinary angles α_i , here $i = 1, \dots, n$.

Suppose that there exist a convex polygon $A_1 A_2 \dots A_n$ with consecutive angles α_i for $i = 1, \dots, n$. Let also the pair of vectors $\overline{A_2 A_3}$ and $\overline{A_2 A_1}$ defines the positive orientation of the plane (otherwise we apply to the polygon $A_1 A_2 \dots A_n$ some \mathcal{L} -affine transformation changing the orientation and come to the initial position).

Let $B_1 = O + \overline{A_n A_1}$, and $B_i = O + \overline{A_{i-1} A_i}$ for $i = 2, \dots, n$. We put by definition

$$\beta_i = \begin{cases} \angle B_i O B_{i+1}, & \text{if } i = 1, \dots, n-1 \\ \angle B_n O B_1, & \text{if } i = n \end{cases}.$$

Consider the union of the sails for all β_i . This lattice oriented broken line is of the class of the extended angle with the normal form $2\pi + 0$. The LSLS-sequence for this broken line contains exactly $n - 1$ elements that are not contained in the LLS-sequences for the sails of β_i . Denote these numbers by m_1, \dots, m_{n-1} , and the set $\{m_1, \dots, m_{n-1}\}$ by M . Then

$$\sum_{M, i=1}^n \overline{\beta_i} = 2\pi.$$

From the definition of β_i for $i = 1, \dots, n$ it follows that $\beta_i \cong \pi - \alpha_i$. Therefore,

$$\sum_{M,i=1}^n \overline{\pi - \alpha_i} = 2\pi.$$

The proof of the first part of the statement is completed.

Suppose now, that there exist a set of integers $M = \{m_1, \dots, m_{n-1}\}$ such that

$$\sum_{M,i=1}^n \overline{\pi - \alpha_i} = 2\pi.$$

This implies that there exist lattice points $B_1 = (1, 0)$, $B_i = (x_i, y_i)$, for $i = 2, \dots, n-1$, and $B_n = (-1, 0)$ such that

$$\angle B_i O B_{i-1} \cong \pi - \alpha_{i-1}, \text{ for } i = 2, \dots, n, \quad \text{and} \quad \angle B_1 O B_n \cong \pi - \alpha_n.$$

Denote by M the lattice point

$$O + \sum_{i=1}^n \overline{OB_i}.$$

Since all α_i are non-zero, the angles $\pi - \alpha_i$ are ordinary. Hence, the origin O is an interior point of the convex hull of the points B_i for $i = 1, \dots, k$. This implies that there exist two consecutive lattice points B_s and B_{s+1} (or B_n and B_1), such that the lattice triangle $\triangle B_s M B_{s+1}$ contains O and the edge $B_s B_{s+1}$ does not contain O . Therefore,

$$O = \lambda_1 \overline{OM} + \lambda_2 \overline{OB_i} + \lambda_3 \overline{OB_{i+1}},$$

where λ_1 is a positive integer, and λ_2 and λ_3 are nonnegative integers. So there exist positive integers a_i , where $i = 1, \dots, n$, such that

$$O = O + \sum_{i=1}^n (a_i \overline{OB_i}).$$

Put by definition $A_0 = O$, and $A_i = A_{i-1} + a_i \overline{OB_i}$ for $i = 2, \dots, n$. The broken line $A_0 A_1 \dots A_n$ is lattice and by the above it is closed (i. e. $A_0 = A_n$). By construction, the ordinary angle at the vertex A_i of the closed lattice broken line is \mathcal{L}_+ -congruent to α_i ($i = 1, \dots, n$). Since the integers a_i are positive for $i = 1, \dots, n$ and the vectors $\overline{OB_i}$ are all in the counterclockwise order, the broken line is a convex polygon.

The proof of Theorem 4.6 is completed.

REMARK 4.7. Theorem 4.6 generalizes the statement of Theorem 2.2a. Note that the direct generalization of Theorem 2.2b is false: the ordinary angles do not uniquely determine the \mathcal{L}_+ -affine homothety types of convex polygons. See an example on Figure 10.



FIGURE 10. An example of different types of polygons with the \mathcal{L}_+ -congruent ordinary angles.

Appendix A. On global relations on algebraic singularities of complex projective toric varieties corresponding to integer-lattice triangles

In this appendix we describe an application of theorems on sums of lattice tangents for the angles of lattice triangles and lattice convex polygons to theory of complex projective toric varieties. We refer the reader to the general definitions of theory of toric varieties to the works of V. I. Danilov [4], G. Ewald [5], W. Fulton [6], and T. Oda [18].

Let us briefly recall the definition of complex projective toric varieties associated to lattice convex polygons. Consider a lattice convex polygon P with vertices A_0, A_1, \dots, A_n . Let the intersection of this (closed) polygon with the lattice consists of the points $B_i = (x_i, y_i)$ for $i = 0, \dots, m$. Let also $B_i = A_i$ for $i = 0, \dots, n$. Denote by Ω the following set in $\mathbb{C}P^m$:

$$\{(t_1^{x_1} t_2^{y_1} t_3^{-x_1-y_1} : t_1^{x_2} t_2^{y_2} t_3^{-x_2-y_2} : \dots : t_1^{x_m} t_2^{y_m} t_3^{-x_m-y_m}) \mid t_1, t_2, t_3 \in \mathbb{C} \setminus \{0\}\}.$$

The closure of the set Ω in the natural topology of $\mathbb{C}P^m$ is called the *complex toric variety associated with the polygon P* and denoted by X_P .

For any $i = 0, \dots, m$ we denote by \tilde{A}_i the point $(0 : \dots : 0 : 1 : 0 : \dots : 0)$ where 1 stands on the $(i+1)$ -th place.

From general theory it follows that:

- a) the set X_P is a complex projective complex-two-dimensional variety with isolated algebraic singularities;
- b) the complex toric projective variety contains the points \tilde{A}_i for $i = 0, \dots, n$ (where $n+1$ is the number of vertices of convex polygon);
- c) the points of $X_P \setminus \{\tilde{A}_0, \tilde{A}_1, \dots, \tilde{A}_n\}$ are non-singular;
- d) the point \tilde{A}_i for any integer i satisfying $0 \leq i \leq n$ is singular iff the corresponding ordinary angle α_i at the vertex A_i of the polygon P is not \mathcal{L} -congruent to $\text{larctan}(1)$;

- e) the algebraic singularity at \tilde{A}_i for any integer i satisfying $0 \leq i \leq n$ is uniquely determined by the \mathcal{L} -affine type of the non-oriented sail of the lattice angle α_i .

The algebraic singularity is said to be *toric* if there exists a projective toric variety with the given algebraic singularity.

Note that the \mathcal{L} -affine classes of non-oriented sails for angles α and β coincide iff $\beta \cong \alpha$, or $\beta \cong \alpha^t$. This allows us to associate to any complex-two-dimensional toric algebraic singularity, corresponding to the sail of the angle α , the unordered pair of rationals (a, b) , where $a = \text{l tan } \alpha$ and $b = \text{l tan } \alpha^t$.

REMARK A.1. Note that the continued fraction for the sail α is slightly different to the Hirzebruch-Jung continued fractions for toric singularities (see the works [9] by H. W. E. Jung, and [8] by F. Hirzebruch). The relations between these continued fractions is described in the paper [19] by P. Popescu-Pampu.

COROLLARY A.2. *Suppose, that we are given by three complex-two-dimensional toric singularities defined by pairs of rationals (a_i, b_i) for $i = 1, 2, 3$. There exist a complex toric variety associated with some triangle with these three singularities iff there exist a permutation $\sigma \in S_3$ and the rationals c_i from the sets $\{a_i, b_i\}$ for $i = 1, 2, 3$, such that the following conditions hold:*

- i) *the continued fraction $]c_{\sigma(1)}, -1, c_{\sigma(2)}[$ is either negative, or greater than $c_{\sigma(1)}$, or equals ∞ ;*
- ii) $]c_{\sigma(1)}, -1, c_{\sigma(2)}, -1, c_{\sigma(3)}[= 0$.

We note again that we use odd continued fractions for c_1, c_2 , and c_3 in the statement of the above proposition (see Subsection 2.1 for the notation of continued fractions).

PROOF. The proposition follows directly from Theorem 2.2a.

PROPOSITION A.3. *For any collection (with multiplicities) of complex-two-dimensional toric algebraic singularities there exist a complex-two-dimensional toric projective variety with exactly the given collection of toric singularities.*

For the proof of Proposition A.3 we need the following lemma.

LEMMA A.4. *For any collection of ordinary angles α_i ($i = 1, \dots, n$), there exist an integer $k \geq n-1$ and a k -tuple of integers $M = (m_1, \dots, m_k)$, such that*

$$\overline{\alpha}_1 +_{m_1} \cdots +_{m_{n-1}} \overline{\alpha}_n +_{m_n} \text{larctan}(1) +_{m_{n+1}} \cdots +_{m_k} \text{larctan}(1) = 2\pi.$$

PROOF. Consider any collection of ordinary angles α_i ($i = 1, \dots, n$) and denote

$$\Phi = \overline{\alpha}_1 +_1 \overline{\alpha}_2 +_1 \cdots +_1 \overline{\alpha}_n.$$

There exist an oriented lattice broken line for the angle Φ with the LSLS-sequence with positive elements. Hence, $\Phi \hat{\cong} \varphi + 0\pi$.

If $\varphi \cong \text{larctan}(1)$, we have

$$\Phi +_{-2} \text{larctan}(1) +_{-2} \text{larctan}(1) +_{-2} \text{larctan}(1) = 2\pi.$$

Then $k = n + 2$, and $M = (1, \dots, 1, -2, -2, -2)$.

Suppose now $\varphi \not\cong \text{larctan}(1)$, then the following holds

$$\bar{\varphi} +_{-1} \overline{\pi - \varphi} +_{-2} \text{larctan}(1) +_{-2} \text{larctan}(1) = 2\pi.$$

Consider the sail for the angle $\pi - \varphi$. Suppose the sequence of all its lattice points (not only vertices) is B_0, \dots, B_s (with the order coinciding with the order of the sail). Then we have

$$\angle B_i O B_{i+1} \hat{\cong} \text{larctan}(1) \quad \text{for any } i = 1, \dots, s.$$

Denote by b_i the values of $\text{lsin } \angle B_i O B_{i+1}$ for $i = 1, \dots, s$. Then we have

$$\begin{aligned} &\bar{\varphi} +_{-2} \text{larctan}(1) +_{-2} \text{larctan}(1) +_{-2} \text{larctan}(1) \\ &= \overline{\alpha_1} +_1 \overline{\alpha_2} +_1 \dots +_1 \overline{\alpha_n} +_{-1} \text{larctan}(1) +_{b_1} \text{larctan}(1) +_{b_2} \\ &\quad \dots +_{b_s} \text{larctan}(1) +_{-2} \text{larctan}(1) +_{-2} \text{larctan}(1) +_{-2} \text{larctan}(1) \\ &= 2\pi. \end{aligned}$$

Therefore, $k = n + s + 3$, and

$$M = \underbrace{(1, 1, \dots, 1, 1)}_{(n-1)\text{-times}}, -1, b_1, \dots, b_s, -2, -2, -2).$$

The proof of Lemma A.4 is completed.

PROOF OF THE STATEMENT OF THE PROPOSITION A.3. Consider an arbitrary collection of two-dimensional toric algebraic singularities. Suppose that they are represented by ordinary angles α_i ($i = 1, \dots, n$). By Lemma A.4 there exist an integer $k \geq n - 1$ and a k -tuple of integers $M = (m_1, \dots, m_k)$, such that

$$\overline{(\pi - \alpha_1)} +_{m_1} \dots +_{m_{n-1}} \overline{(\pi - \alpha_n)} +_{m_n} \text{larctan}(1) +_{m_{n+1}} \dots +_{m_k} \text{larctan}(1) = 2\pi.$$

By Theorem 4.6 there exist a convex polygon $P = A_0 \dots A_k$ with angles \mathcal{L}_+ -congruent to the ordinary angles α_i ($i = 1, \dots, n$), and $k - n + 1$ angles $\text{larctan}(1)$.

By the above, the toric variety X_P is nonsingular at points of $P_X \setminus \{\tilde{A}_0, \tilde{A}_1, \dots, \tilde{A}_k\}$. It is also nonsingular at the points \tilde{A}_i with the corresponding ordinary

angles \mathcal{L} -congruent to $\text{larctan}(1)$. The collection of the toric singularities at the remaining points coincide with the given collection.

This concludes the proof of Proposition A.3.

On Figure 11 we show an example of the polygon for a projective toric variety with the unique toric singularity, represented by the sail of $\text{larctan}(7/5)$. The ordinary angle α on the figure is \mathcal{L}_+ -congruent to $\text{larctan}(7/5)$, the angles β and γ are \mathcal{L}_+ -congruent to $\text{larctan}(1)$.




FIGURE 11. Constructing a polygon with all angles \mathcal{L}_+ -congruent to $\text{larctan}(1)$ except one angle that is \mathcal{L}_+ -congruent to $\text{larctan}(7/5)$.

Appendix B. On \mathcal{L} -congruence criterions for lattice triangles

Here we discuss the \mathcal{L} -congruence criterions for lattice triangles. By the first criterion of \mathcal{L} -congruence for lattice triangles we obtain that the number of \mathcal{L} -congruence classes for lattice triangles with bounded lattice area is finite. We write down the numbers of \mathcal{L} -congruence classes for triangles with lattice area less then or equal to 20.

On criterions of lattice triangle \mathcal{L} -congruence. We start with the study of lattice analogs for the first, the second, and the third Euclidean criterions of triangle congruence.

STATEMENT B.1 (The first criterion of lattice triangle \mathcal{L} -congruence). *Consider two lattice triangles $\triangle ABC$ and $\triangle A'B'C'$. Suppose that the edge AB is \mathcal{L} -congruent to the edge $A'B'$, the edge AC is \mathcal{L} -congruent to the edge $A'C'$, and the ordinary angle $\angle CAB$ is \mathcal{L} -congruent to the ordinary angle $\angle C'A'B'$, then the triangle $\triangle A'B'C'$ is \mathcal{L} -congruent to the $\triangle ABC$.*

It turns out that the second and the third criterions taken from Euclidean geometry do not hold. The following two examples illustrate these phenomena.

EXAMPLE B.2. The second criterion of triangle \mathcal{L} -congruence does not hold in lattice geometry. On Figure 12 we show two lattice triangles $\triangle ABC$ and $\triangle A'B'C'$. The edge AB is \mathcal{L} -congruent to the edge $A'B'$ (here $l\ell(A'B') = l\ell(AB) = 4$). The ordinary angle $\angle ABC$ is \mathcal{L} -congruent to the ordinary angle $\angle A'B'C'$ (since $\angle ABC \cong \angle A'B'C' \cong \arctan(1)$), and the ordinary angle $\angle CAB$ is \mathcal{L} -congruent to the ordinary angle $\angle C'A'B'$ (since $\angle CAB \cong \angle C'A'B' \cong \arctan(1)$). The triangle $\triangle A'B'C'$ is not \mathcal{L} -congruent to the triangle $\triangle ABC$, since $IS(\triangle ABC) = 4$ and $IS(\triangle A'B'C') = 8$.



FIGURE 12. The second criterion of triangle \mathcal{L} -congruence does not hold.

EXAMPLE B.3. The third criterion of triangle \mathcal{L} -congruence does not hold in lattice geometry. On Figure 13 we show two lattice triangles $\triangle ABC$ and $\triangle A'B'C'$. All edges of both triangles are \mathcal{L} -congruent (of length one), but the triangles are not \mathcal{L} -congruent, since $IS(\triangle ABC) = 1$ and $IS(\triangle A'B'C') = 3$.




FIGURE 13. The third criterion of triangle \mathcal{L} -congruence does not hold.

Instead of the second and the third criteria there exists the following additional criterion of lattice triangles \mathcal{L} -congruence.

STATEMENT B.4 (An additional criterion of lattice triangle integer-congruence). *Consider two lattice triangles $\triangle ABC$ and $\triangle A'B'C'$ of the same lattice area. Suppose that the ordinary angle $\angle ABC$ is \mathcal{L} -congruent to the ordinary angle $\angle A'B'C'$, the ordinary angle $\angle CAB$ is \mathcal{L} -congruent to the ordinary angle $\angle C'A'B'$, the ordinary angle $\angle BCA$ is \mathcal{L} -congruent to the ordinary angle $\angle B'C'A'$, then the triangle $\triangle A'B'C'$ is \mathcal{L} -congruent to the triangle $\triangle ABC$.*

In the following example we show that the additional criterion of lattice triangle \mathcal{L} -congruence is not improvable.

EXAMPLE B.5. On Figure 14 we show an example of two lattice non-equivalent triangles $\triangle ABC$ and $\triangle A'B'C'$ of the same lattice area equals 4

and the same ordinary angles $\angle ABC$, $\angle CAB$, and $\angle A'B'C'$, $\angle C'A'B'$ all \mathcal{L} -equivalent to the angle $\arctan(1)$, but $\triangle ABC \not\cong \triangle A'B'C'$.




FIGURE 14. The additional criterion of lattice triangle \mathcal{L} -congruence is not improvable.

Lattice triangles of small area. The above criterions allows to enumerate all lattice triangles of small lattice area up to the lattice equivalence. In the following table we write down the numbers $N(d)$ of nonequivalent lattice triangles of lattice area d for $d \leq 20$.

d	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
$N(d)$	1	1	2	3	2	4	4	5	5	6	4	10	6	8	8	11	6	13	8	14

As it is easy to show, we always have $d/3 \leq N(d) \leq d$. The asymptotic behaviour of $N(d)$ and even of the average of $N(d)$ (if they exist) is unknown to the author.

Appendix C. Some unsolved question on lattice trigonometry

We conclude this paper with a small collection of unsolved questions.

Let us start with some questions on elementary definitions of lattice trigonometry. In this paper we do not show any geometrical meaning of lattice cosine. Here arise the following question.

PROBLEM 1. Find a natural description of lattice cosine for ordinary angles in terms of lattice invariants of the corresponding sublattices.

This problem seems to be close to the following one.

PROBLEM 2. Does there exist a lattice analog of the cosine formula for the angles of triangles in Euclidean geometry?

Let us continue with questions on lattice analogs of classical trigonometric formulas for trigonometric functions of angles of triangles in Euclidean geometry.

PROBLEM 3. a) Knowing the lattice trigonometric functions for lattice angles α , β and integer n , find the explicit formula for the lattice trigonometric functions of the extended angle $\bar{\alpha} +_n \bar{\beta}$.

b) Knowing the lattice trigonometric functions for a lattice angle α , an integer m , and positive integer m , find the explicit formula for the lattice trigonometric functions of the extended angle

$$\sum_{M, i=1}^l \bar{\alpha},$$

where $M = (m, \dots, m)$ is an n -tuple.

Now we formulate a problem on generalization of the statement of Theorem 2.2b to the case of n ordinary angles. Such generalization is important in toric geometry and theory of multidimensional continued fractions.

PROBLEM 4. Find a necessary and sufficient conditions for the existence of an n -gon with the given ordered sequence of ordinary angles $(\alpha_1, \dots, \alpha_n)$ and the consistent sequence of lattice lengths of the edges (l_1, \dots, l_n) in terms of continued fractions for $n \geq 4$.

We conclude this paper with the following problem. We remind that $(N(d))$ is the numbers of nonequivalent lattice triangles having the lattice area being equal to d (see Appendix B).

PROBLEM 5. Find an explicit formula for the numbers $N(d)$.

REFERENCES

1. Arnold, V. I., *Continued fractions*, M., Moscow Center of Continuous Mathematical Education, 2002.
2. Arnold, V. I., *Statistics of integer convex polygons*, *Funct. Anal. Appl.* 14(2) (1980), 1–3.
3. Bárány, I., Vershik, A. M., *On the number of convex lattice polytopes*, *Geom. Funct. Anal.* 2(4) (1992), 381–393.
4. Danilov, V. I., *The geometry of toric varieties*, *Uspekhi Mat. Nauk.* 33(2) (1978), 85–134.
5. Ewald, G., *Combinatorial Convexity and Algebraic Geometry*, *Grad. Texts in Math.* 168 (1996).
6. Fulton, W., *Introduction to Toric Varieties*, *Ann. of Math. Stud.* 131 (1993).
7. Hinchin, A. Ya., *Continued Fractions*, M.: FISMATGIS, 1961.
8. Hirzebruch, F., *Über vierdimensionale Riemannsche Flächen Mehrdeutiger analytischer Funktionen von zwei komplexen Veränderlichen*, *Math. Ann.* 126 (1953), 1–22.
9. Jung, H. W. E., *Darstellung der Funktionen eines algebraischen Körpers zweier unabhängigen Veränderlichen x, y in der Umgebung einer Stelle $x = a, y = b$* , *J. Reine Angew. Math.* 133 (1908), 289–314.
10. Karpenkov, O., *On tori decompositions associated with two-dimensional continued fractions of cubic irrationalities*, *Funct. Anal. Appl.* 38(2) (2004), 28–37.
11. Karpenkov, O., *Classification of lattice-regular lattice convex polytopes*, <http://arxiv.org/abs/math.CO/0602193>.

12. Khovanskii, A. G., Pukhlikov, A., *Finitely additive measures of virtual polytopes*, Algebra and Analysis, 4(2) (1992), 161–185; translation in St. Petersburg Math. J. 4(2) (1993), 337–356.
13. Khovanskii, A. G., Pukhlikov, A., *A Riemann-Roch theorem for integrals and sums of quasi-polynomials over virtual polytopes*, Algebra and Analysis 4(4) (1992), 188–216; translation in St. Petersburg Math. J. 4(4) (1993), 789–812.
14. Klein, F., *Ueber eine geometrische Auffassung der gewöhnliche Kettenbruchentwicklung*, Nachr. Ges. Wiss. Göttingen Math-Phys. Kl. 3 (1891), 357–359.
15. Kontsevich, M. L., and Suhov, Yu. M., *Statistics of Klein Polyhedra and Multidimensional Continued Fractions*, Amer. Math. Soc. Transl. 197(2) (1999), 9–27.
16. Korkina, E. I., *Two-dimensional continued fractions. The simplest examples*, Proc. Steklov Inst. Math. 209 (1995), 243–166.
17. Lachaud, G., *Voiles et Polyèdres de Klein*, preprint n 95-22, Laboratoire de Mathématiques Discrètes du C.N.R.S., Luminy (1995).
18. Oda, T., *Convex bodies and Algebraic Geometry, An Introduction to the Theory of Toric Varieties*, Ergeb. Math. Grenzgeb. 145 (1988)
19. Popescu-Pampu, P., *The Geometry of Continued Fractions and The Topology of Surface Singularities*, ArXiv:math.GT/0506432, v1 (2005).

CEREMADE - UMR 7534
UNIVERSITÉ PARIS-DAUPHINE FRANCE
75775 PARIS CEDEX 16
FRANCE
E-mail: karpenk@mccme.ru