

# A CLASSIFICATION OF $\mathbb{C}$ -FUCHSIAN SUBGROUPS OF PICARD MODULAR GROUPS

JOUNI PARKKONEN and FRÉDÉRIC PAULIN

## Abstract

Given an imaginary quadratic extension  $K$  of  $\mathbb{Q}$ , we give a classification of the maximal nonelementary subgroups of the Picard modular group  $\mathrm{PSU}_{1,2}(\mathcal{O}_K)$  preserving a complex geodesic in the complex hyperbolic plane  $\mathbb{H}_{\mathbb{C}}^2$ . Complementing work of Holzapfel, Chinburg-Stover and Möller-Toledo, we show that these maximal  $\mathbb{C}$ -Fuchsian subgroups are arithmetic, arising from a quaternion algebra  $(\frac{D, D_K}{\mathbb{Q}})$  for some explicit  $D \in \mathbb{N} - \{0\}$  and  $D_K$  the discriminant of  $K$ . We thus prove the existence of infinitely many orbits of  $K$ -arithmetic chains in the hypersphere of  $\mathbb{P}_2(\mathbb{C})$ .

## 1. Introduction

Let  $h$  be a Hermitian form with signature  $(1, 2)$  on  $\mathbb{C}^3$ . The projective special unitary Lie group  $\mathrm{PSU}_h$  of  $h$  contains exactly two conjugacy classes of connected Lie subgroups locally isomorphic to  $\mathrm{SL}_2(\mathbb{R})$ . The subgroups in one class, isomorphic to  $\mathrm{SU}(1, 1) \simeq \mathrm{SL}_2(\mathbb{R})$ , preserve a complex projective line for the projective action of  $\mathrm{PSU}_h$  on the projective plane  $\mathbb{P}_2(\mathbb{C})$ , and those of the other class, isomorphic to  $\mathrm{SO}_0(2, 1) \simeq \mathrm{PSL}_2(\mathbb{R})$ , preserve a totally real subspace. The groups  $\mathrm{PSL}_2(\mathbb{R})$  and  $\mathrm{PSU}_h$  act as the groups of holomorphic isometries, respectively, on the upper halfplane model  $\mathbb{H}_{\mathbb{R}}^2$  of the real hyperbolic space and on the projective model  $\mathbb{H}_{\mathbb{C}}^2$  of the complex hyperbolic plane defined using the form  $h$ . If  $\Gamma$  is a discrete subgroup of  $\mathrm{PSU}_h$ , the intersections of  $\Gamma$  with the connected Lie subgroups locally isomorphic to  $\mathrm{SL}_2(\mathbb{R})$  are its *Fuchsian subgroups*, and the Fuchsian subgroups preserving a complex projective line are called  *$\mathbb{C}$ -Fuchsian subgroups*. We refer to Section 2 for more precise definitions and comments on the terminology.

Let  $K$  be an imaginary quadratic number field, with discriminant  $D_K$  and ring of integers  $\mathcal{O}_K$ . We consider the Hermitian form  $h$  defined by

$$(z_0, z_1, z_2) \mapsto -z_0 \overline{z_2} - z_2 \overline{z_0} + z_1 \overline{z_1}.$$

The *Picard modular group*  $\Gamma_K = \mathrm{PSU}_h(\mathcal{O}_K)$  is a nonuniform arithmetic lattice of  $\mathrm{PSU}_h$ , see for instance [13, Chap. 5] and subsequent works of Falbel, Parker,

Francis, Lax, Xie-Wang-Jiang, and many others for information on these groups. In this paper, we classify the maximal  $\mathbb{C}$ -Fuchsian subgroups of  $\Gamma_K$ , and we explicit their arithmetic structures. The results stated in this introduction do not depend on the choice of the Hermitian form  $h$  of signature  $(2, 1)$  defined over  $K$ , since the algebraic groups over  $\mathbb{Q}$  whose groups of  $\mathbb{Q}$  points are  $\mathrm{PSU}_h(K)$  depend up to commensurability only on  $K$  and not on  $h$ , see for instance [26, § 3.1].

When  $G = \mathrm{PSL}_2(\mathbb{C})$ , there is exactly one conjugacy class of connected Lie subgroups of  $G$  isomorphic to  $\mathrm{PSL}_2(\mathbb{R})$ . When  $\Gamma$  is the *Bianchi group*  $\mathrm{PSL}_2(\mathcal{O}_K)$ , the classification analogous to the one we describe is due to MacLachlan and Reid (see [17], [18] and [19, Chap. 9]). They proved that the maximal nonelementary Fuchsian subgroups of  $\mathrm{PSL}_2(\mathcal{O}_K)$  are commensurable up to conjugacy in  $\mathrm{PSL}_2(\mathbb{C})$  with the stabilisers of the circles  $|z|^2 = D$  for  $D \in \mathbb{N} - \{0\}$ , when  $\mathrm{PSL}_2(\mathbb{C})$  acts projectively (by homographies) on the projective line  $\mathbb{P}_1(\mathbb{C}) = \mathbb{C} \cup \{\infty\}$ , and that all these subgroups arise from explicit quaternion algebras over  $\mathbb{Q}$ . For information on Bianchi groups, see for instance [9] and the references of [18].

More generally, given a connected semisimple real Lie group  $G$  with finite center and without compact factors, there is a nonempty finite set of infinite conjugacy classes of connected Lie subgroups of  $G$  locally isomorphic to  $\mathrm{SL}_2(\mathbb{R})$ , unless  $G$  itself is locally isomorphic to  $\mathrm{SL}_2(\mathbb{R})$ . The structure of the set of these subgroups plays an important role for the classification of the linear representations of  $G$ , and for the classification of the groups  $G$  themselves, see for instance [15], [25] among others. Given a discrete subgroup  $\Gamma$  of  $G$ , it is again interesting to study the *Fuchsian subgroups* of  $\Gamma$ , that is, the intersections of  $\Gamma$  with these Lie subgroups, to classify the maximal ones and to see, when  $\Gamma$  is arithmetic, if its maximal Fuchsian subgroups are also arithmetic (see Proposition 3.1 for a positive answer) with an explicit arithmetic structure. From now on,  $G = \mathrm{PSU}_h$ .

We first prove (see Proposition 3.2 and just after) that a nonelementary  $\mathbb{C}$ -Fuchsian subgroup  $\Gamma'$  of  $\Gamma_K$  preserves a unique projective point  $[z_0 : z_1 : z_2]$  with  $z_0, z_1, z_2$  relatively prime in  $\mathcal{O}_K$ . We define the *discriminant* of  $\Gamma'$  as  $\Delta_{\Gamma'} = h(z_0, z_1, z_2)$ . For any positive integer  $D$ , let

$$\Gamma_{K,D} = \mathrm{Stab}_{\Gamma_K}[-D : 0 : 1].$$

In Section 3, we prove the following classification result (see [18, Thm. 1] and [19, Thm. 9.6.2] in the Bianchi group case).

**THEOREM 1.1.** *Let  $D \in \mathbb{N} - \{0\}$ . The set of  $\Gamma_K$ -conjugacy classes of maximal nonelementary  $\mathbb{C}$ -Fuchsian subgroups of  $\Gamma_K$  with discriminant  $D$  is finite*

and nonzero. Every maximal nonelementary  $\mathbb{C}$ -Fuchsian subgroup of  $\Gamma_K$  with discriminant  $D$  is commensurable up to conjugacy in  $\text{PSU}_h$  with  $\Gamma_{K,2D}$ .

In the course of the proof of this result (see Lemma 3.4 and Corollary 4.2), we prove a criterion for when two groups  $\Gamma_{K,D}$  for  $D \in \mathbb{N} - \{0\}$  are commensurable up to conjugacy in  $\text{PSU}_h$ . A further application of this condition shows that every maximal nonelementary  $\mathbb{C}$ -Fuchsian subgroup of  $\Gamma_K$  is commensurable up to conjugacy in  $\text{PSU}_h$  with  $\Gamma_{K,D}$  for a squarefree positive integer  $D$ .


Recall (see for instance [10]) that a *chain*<sup>1</sup> is the intersection of the *Poincaré hypersphere*

$$\mathcal{HS} = \{[z] \in \mathbb{P}_2(\mathbb{C}) : h(z) = 0\}$$

with a complex projective line (if nonempty and not a singleton). It is *K-arithmetic* if its stabiliser in  $\Gamma_K$  has a dense orbit in it (see Section 3 for an explanation of the terminology).

**COROLLARY 1.2.** *There are infinitely many  $\Gamma_K$ -orbits of K-arithmetic chains in the hypersphere  $\mathcal{HS}$ .*

The figure below shows part of the image under vertical projection in the Heisenberg group of the orbit under  $\Gamma_K$  of a *K*-arithmetic chain whose stabiliser has discriminant 10, when  $K = \mathbb{Q}(i)$ .



We say that a subgroup of  $\text{PSU}_h$  arises from a quaternion algebra  $A$  defined over  $\mathbb{Q}$  if it is commensurable in  $\text{PSU}_h$  with  $\sigma(A(\mathbb{Z})^1)$  for some  $\mathbb{Q}$ -algebra morphism  $\sigma : A \rightarrow \mathcal{M}_3(\mathbb{C})$ , where  $A(\mathbb{Z})^1$  is the group of norm 1 elements in

---

<sup>1</sup> a notion attributed to von Staudt in [3, footnote 3]

$A(\mathbb{Z})$ . In Section 4, we prove the following result (see [19, Thm. 9.6.3] in the Bianchi group case).

**THEOREM 1.3.** *Every nonelementary  $\mathbb{C}$ -Fuchsian subgroup of  $\Gamma_K$  of discriminant  $D$  is conjugate in  $\mathrm{PSU}_h$  to a subgroup of  $\mathrm{PSU}_h$  arising from the quaternion algebra  $(\frac{D, D_K}{\mathbb{Q}})$ .*

The classification of the quaternion algebras over  $\mathbb{Q}$  then allows to classify, up to commensurability and conjugacy in  $\mathrm{PSU}_h$ , the maximal nonelementary  $\mathbb{C}$ -Fuchsian subgroups of  $\Gamma_K$ : two such groups, with discriminant  $D$  and  $D'$  are commensurable up to conjugacy in  $\mathrm{PSU}_h$  if and only if the quaternion algebras  $(\frac{D, D_K}{\mathbb{Q}})$  and  $(\frac{D', D_K}{\mathbb{Q}})$  are isomorphic (see Corollary 4.2). This holds for instance if and only if the quadratic forms  $D_K x^2 + D y^2 - D D_K z^2$  and  $D_K x^2 + D' y^2 - D' D_K z^2$  are equivalent over  $\mathbb{Q}$  (see for instance [19, Coro. 2.3.5]).

We conclude the introduction with a brief review of previous work related to this paper. The terminology of this paragraph will not be needed in this paper, and we refer to the quoted references for details. The existence of a bijection between wide commensurability classes of  $\mathbb{C}$ -Fuchsian subgroups of  $\Gamma_K$  and isomorphism classes of quaternion algebras over  $\mathbb{Q}$  unramified at infinity and ramified at all finite places which do not split in  $K/\mathbb{Q}$  is a particular case of a 2011 result of Chinburg-Stover (see Theorem 2.2 in version 3 of [4] and [5, Theo. 4.1]), which proves such a result for all arithmetic lattices of simple type (which include the Picard modular groups) in  $\mathrm{SU}_{2,1}$ . In particular, the existence of this bijection (and our Corollary 4.4) should be attributed to Chinburg-Stover (although they say it was known by experts). Möller-Toledo in [20] also give a description of the quotients by the maximal  $\mathbb{C}$ -Fuchsian subgroups of the real hyperbolic planes they preserve, and more generally of all Shimura curves in Shimura surfaces of the first type (which include the complex surfaces  $\Gamma_K \backslash \mathbb{H}_{\mathbb{C}}^2$ ). We believe that our precise correspondence brings interesting effective and geometric information to the picture.

**ACKNOWLEDGEMENTS.** The first author thanks the Väisälä foundation and the FIM of ETH Zürich for their support during the preparation of this paper. The second author thanks the Väisälä foundation and its financial support for a fruitful visit to the University of Jyväskylä and the Nordic snows. This work is supported by the NSF Grant no. 093207800, while the second author was in residence at the MSRI, Berkeley CA, during the Spring 2015 semester. We thank Y. Benoist and M. Burger for interesting discussions on this paper. We warmly thank M. Stover for informing us (after we posted a first version of this paper on ArXiv) about the third version on ArXiv of the paper [4] and many other references, including [20]. We thank the referee for many very helpful remarks and suggestions which improved this paper significantly.

## 2. The complex hyperbolic plane

Let  $h$  be the nondegenerate Hermitian form

$$h(z) = -z_0 \bar{z}_2 - z_2 \bar{z}_0 + |z_1|^2 = -2 \operatorname{Re}(z_0 \bar{z}_2) + |z_1|^2$$

of signature  $(1, 2)$  on  $\mathbb{C}^3$  with coordinates  $(z_0, z_1, z_2)$ , and let  $\langle \cdot, \cdot \rangle$  be the associated Hermitian product. The point  $z = (z_0, z_1, z_2) \in \mathbb{C}^3$  and the corresponding element  $[z] = [z_0 : z_1 : z_2] \in \mathbb{P}_2(\mathbb{C})$  (using homogeneous coordinates) is *negative*, *null* or *positive* according to whether  $h(z) < 0$ ,  $h(z) = 0$  or  $h(z) > 0$ . The *negative/null/positive cone* of  $h$  is the subset of negative/null/positive elements of  $\mathbb{P}_2(\mathbb{C})$ .

The negative cone of  $h$  endowed with the distance  $d$  defined by

$$\cosh^2 d([z], [w]) = \frac{|\langle z, w \rangle|^2}{h(z)h(w)}$$

is the *complex hyperbolic plane*  $\mathbb{H}_{\mathbb{C}}^2$ . The distance  $d$  is the distance of a Riemannian metric with pinched negative sectional curvature  $-4 \leq K \leq -1$ . The linear action of the special unitary group of  $h$

$$\mathrm{SU}_h = \{g \in \mathrm{SL}_3(\mathbb{C}) : h \circ g = h\}$$

on  $\mathbb{C}^3$  (where  $h \circ g : z \mapsto h(gz)$ ) induces a projective action on  $\mathbb{P}_2(\mathbb{C})$  with kernel  $\mathbb{U}_3 \operatorname{Id}$ , where  $\mathbb{U}_3$  is the group of third roots of unity. This projective action preserves the negative, null and positive cones of  $h$  in  $\mathbb{P}_2(\mathbb{C})$ , and is transitive on each of them. The restriction to  $\mathbb{H}_{\mathbb{C}}^2$  of the quotient group  $\mathrm{PSU}_h = \mathrm{SU}_h / (\mathbb{U}_3 \operatorname{Id})$  of  $\mathrm{SU}_h$  is the holomorphic isometry group of  $\mathbb{H}_{\mathbb{C}}^2$ . Note that the inclusion  $\mathrm{SU}_h \rightarrow \mathrm{U}_h$  induces a Lie group isomorphism  $\mathrm{PSU}_h \rightarrow \mathrm{PU}_h$ .

The null cone of  $h$  is the *Poincaré hypersphere*  $\mathcal{HS}$ , which is naturally identified with the boundary at infinity of  $\mathbb{H}_{\mathbb{C}}^2$ . The *Heisenberg group*

$$\mathrm{Heis}_3 = \{[w_0 : w : 1] \in \mathbb{C} \times \mathbb{C} : 2 \operatorname{Re} w_0 = |w|^2\}$$

acts isometrically on  $\mathbb{H}_{\mathbb{C}}^2$  and simply transitively on  $\mathcal{HS} - \{[1 : 0 : 0]\}$  by the action induced by the matrix representation

$$[w_0 : w : 1] \mapsto \begin{pmatrix} 1 & \bar{w} & w_0 \\ 0 & 1 & w \\ 0 & 0 & 1 \end{pmatrix}$$

of  $\mathrm{Heis}_3$  in  $\mathrm{SU}_h$ . The projective transformations induced by these matrices are called *Heisenberg translations*.

If a complex projective line meets  $\mathbb{H}_{\mathbb{C}}^2$ , its intersection with  $\mathbb{H}_{\mathbb{C}}^2$  is a totally geodesic submanifold of  $\mathbb{H}_{\mathbb{C}}^2$ , called a *complex geodesic*. The intersection of

a complex projective line in  $\mathbb{P}_2(\mathbb{C})$  with the Poincaré hypersphere is called a *chain*, if nonempty and not reduced to a point. Each complex projective line  $L$  in  $\mathbb{P}_2(\mathbb{C})$  meeting  $\mathbb{H}_{\mathbb{C}}^2$  (or its associated complex geodesic  $L \cap \mathbb{H}_{\mathbb{C}}^2$ , or its associated chain  $L \cap \mathcal{HS}$ ) is *polar* to a unique positive point  $P_L \in \mathbb{P}_2(\mathbb{C})$ , that is,  $\langle z, P_L \rangle = 0$  for all  $z \in L$  (or equivalently  $z \in L \cap \mathbb{H}_{\mathbb{C}}^2$  or  $z \in L \cap \mathcal{HS}$ ). This element  $P_L$  is the *polar point* of the projective line  $L$ , of the complex geodesic  $L \cap \mathbb{H}_{\mathbb{C}}^2$  and of the chain  $L \cap \mathcal{HS}$ . Conversely, for each positive point  $P$ , there is a unique complex projective line  $P^\perp$  polar to  $P$ , the *polar line* of  $P$ . The intersection of  $P^\perp$  with  $\mathbb{H}_{\mathbb{C}}^2$  is a complex geodesic.

An easy computation (using for instance Equation (42) in [22]) shows that

$$\text{Stab}_{\text{SU}_h}[0 : 1 : 0] = \left\{ \begin{pmatrix} \zeta a & 0 & i\zeta b \\ 0 & \zeta^{-2} & 0 \\ -i\zeta c & 0 & \zeta d \end{pmatrix} : a, b, c, d \in \mathbb{R}, \zeta \in \mathbb{C} \right\}. \quad (1)$$

In particular,  $\text{Stab}_{\text{SU}_h}[0 : 1 : 0]$  is isomorphic to  $(\mathbb{S}^1 \times \text{SL}_2(\mathbb{R}))/\{\pm(1, \text{id})\}$ . It injects in  $\text{PSU}_h$  by the canonical projection  $\text{SU}_h \rightarrow \text{PSU}_h$ . Hence the group  $\text{Stab}_{\text{PSU}_h}[0 : 1 : 0]$  is also isomorphic to  $(\mathbb{S}^1 \times \text{SL}_2(\mathbb{R}))/\{\pm(1, \text{id})\}$ . More generally, by conjugation, if  $P = [z_0 : z_1 : z_2]$  is a positive point in  $\mathbb{P}_2(\mathbb{C})$ , then the stabilizer of  $P$  in  $\text{PSU}_h$  is the almost direct product  $A_P B_P$ , where  $A_P$  is the unique Lie group embedding of  $\text{SL}_2(\mathbb{R})$  in  $\text{PSU}_h$  preserving the complex geodesic polar to  $P$ , and  $B_P$  is the group of complex reflections with fixed point set the projective line polar to  $P$ . The group  $B_P$  is isomorphic to  $\mathbb{S}^1$  and centralizes  $A_P$ . We have  $B_P \cap A_P = \{\pm 1\}$ , and  $A_P$  acts on the normal bundle of the complex geodesic polar to  $P$  either by parallel transport or by its opposite.

The polar chain of  $P$  is

$$C_P = \{[w_0 : w_1 : w_2] \in \mathbb{P}_2(\mathbb{C}) : h(w_0, w_1, w_2) = \langle (w_0, w_1, w_2), (z_0, z_1, z_2) \rangle = 0\},$$

that is  $C_P \cap \text{Heis}_3$  is the set of  $[w_0 : w : 1] \in \text{Heis}_3$  satisfying the equation

$$\left( \frac{|w|^2}{2} + i \text{Im } w_0 \right) \bar{z}_2 - w \bar{z}_1 + \bar{z}_0 = 0.$$

When  $z_2 \neq 0$ , in the coordinates  $(w, 2 \text{Im } w_0) \in \mathbb{C} \times \mathbb{R}$  of  $[w_0 : w : 1] \in \text{Heis}_3$ , this is the equation of an ellipse, whose image under the *vertical projection*  $[w_0 : w : 1] \mapsto w$  is the circle with center  $\frac{\bar{z}_1}{z_2}$  and radius  $\frac{\sqrt{h(z_0, z_1, z_2)}}{|z_2|}$  in  $\mathbb{C}$  given by the equation

$$|w|^2 - 2 \text{Re} \left( w \frac{\bar{z}_1}{z_2} \right) + 2 \text{Re} \left( \frac{\bar{z}_0}{z_2} \right) = 0.$$

If  $z_2 = 0$ , then  $C_P \cap \text{Heis}_3$  is the vertical affine line over  $\frac{\bar{z}_0}{z_1}$ .

We refer to Goldman [10, p. 67] and Parker [21] for the basic properties of  $\mathbb{H}_{\mathbb{C}}^2$ . These references use different Hermitian forms of signature  $(1, 2)$  to define the complex hyperbolic plane, and the curvature is often normalised differently from our definitions. Our choices are consistent with [22] and [23].

### 3. Classification of $\mathbb{C}$ -Fuchsian subgroups of $\Gamma_K$

Before starting to study Fuchsian subgroups of discrete subgroups of  $\mathrm{PSU}_h$ , let us mention that it is a very general fact that the maximal *nonelementary* (that is, not virtually cyclic) Fuchsian subgroups of arithmetic subgroups of  $\mathrm{PSU}_h$  are automatically (arithmetic) lattices of the copy of  $\mathrm{PSL}_2(\mathbb{R})$  containing them.

**PROPOSITION 3.1.** *Let  $G$  be a semisimple connected real Lie group with finite center and without compact factors, and let  $\Gamma$  be a maximal nonelementary Fuchsian subgroup of an arithmetic subgroup  $\tilde{\Gamma}$  of  $G$ . Then  $\Gamma$  is an arithmetic lattice in the copy of the connected Lie group locally isomorphic to  $\mathrm{SL}_2(\mathbb{R})$  containing it.*

**PROOF.** We refer to [29, §3.1] for an elementary introduction to algebraic groups and their Zariski topology. Let  $\underline{G}$  be a semisimple connected algebraic group defined over  $\mathbb{Q}$ , let  $\underline{H}$  be a connected algebraic subgroup of  $\underline{G}$  defined over  $\mathbb{R}$  locally isomorphic to  $\mathrm{SL}_2$ , and assume that  $\Gamma = \underline{H}(\mathbb{R}) \cap \underline{G}(\mathbb{Z})$  is nonelementary in  $\underline{H}(\mathbb{R})$ . As a nonelementary subgroup of a group locally isomorphic to  $\mathrm{SL}_2$  is Zariski-dense in it, and as the Zariski-closure of a subgroup of  $\underline{G}(\mathbb{Z})$  is defined over  $\mathbb{Q}$  (see for instance [29, Prop. 3.1.8]), we hence have that  $\underline{H}$  is defined over  $\mathbb{Q}$ . Therefore by the Borel-Harish-Chandra theorem [2, Thm. 7.8],  $\Gamma = \underline{H}(\mathbb{Z})$  is an arithmetic lattice in  $\underline{H}(\mathbb{R})$ . Since the copies of connected Lie subgroups of  $G$  locally isomorphic to  $\mathrm{SL}_2(\mathbb{R})$  are algebraic (see for instance [29, Prop. 3.1.6]), the result follows.

One of the main points of the rest of the paper will be to determine explicitly the arithmetic structure of  $\Gamma$ , that is the  $\mathbb{Q}$ -structure thus constructed on the group locally isomorphic to  $\mathrm{SL}_2(\mathbb{R})$  containing it, relating it to the arithmetic structure of  $\tilde{\Gamma}$ , that is the given  $\mathbb{Q}$ -structure on  $G$ .

Let  $K$  be an imaginary quadratic number field, with  $D_K$  its discriminant,  $\mathcal{O}_K$  its ring of integers,  $\mathrm{tr} : z \mapsto z + \bar{z}$  its trace and  $N : z \mapsto |z|^2 = z\bar{z}$  its norm. Recall (see for instance [24]) that there exists a squarefree positive integer  $d$  such that  $K = \mathbb{Q}(i\sqrt{d})$ , that  $D_K = -d$  and  $\mathcal{O}_K = \mathbb{Z}\left[\frac{1+i\sqrt{d}}{2}\right]$  if  $d \equiv -1 \pmod{4}$ , and that  $D_K = -4d$  and  $\mathcal{O}_K = \mathbb{Z}[i\sqrt{d}]$  otherwise. Note that  $\mathcal{O}_K$  is stable by conjugation, and that  $\mathrm{tr}$  and  $N$  take integral values on  $\mathcal{O}_K$ . A unit  $x$  in  $\mathcal{O}_K$  is an invertible element in  $\mathcal{O}_K$ . Since  $N : K^\times \rightarrow \mathbb{R}^\times$  is a group morphism, we have  $N(x) = 1$  for every unit  $x$  in  $\mathcal{O}_K$ .

The *Picard modular group* of  $K$ , that we denote by  $\Gamma_K = \text{PSU}_h(\mathcal{O}_K)$ , consists of the images in  $\text{PSU}_h$  of matrices of  $\text{SU}_h$  with entries in  $\mathcal{O}_K$ . It is a nonuniform arithmetic lattice by the result of Borel and Harish-Chandra cited above. Note that every nonuniform arithmetic lattice in  $\text{PSU}_h$  is commensurable to a Picard modular group (see for instance [26, § 3.1]).

A discrete subgroup  $\Gamma$  of  $\text{PSU}_h$  is an *extended  $\mathbb{C}$ -Fuchsian subgroup* if it satisfies one of the following equivalent conditions

- (1)  $\Gamma$  preserves a complex projective line of  $\mathbb{P}_2(\mathbb{C})$  meeting  $\mathbb{H}_{\mathbb{C}}^2$ ,
- (2)  $\Gamma$  fixes a positive point in  $\mathbb{P}_2(\mathbb{C})$ ,
- (3)  $\Gamma$  preserves a chain.

Many references, see for example [7], do not use the word “extended”. But as defined in the introduction, in this paper, a  *$\mathbb{C}$ -Fuchsian subgroup* is a discrete subgroup of  $\text{PSU}_h$  preserving a complex geodesic in  $\mathbb{H}_{\mathbb{C}}^2$  and inducing the parallel transport or its opposite on its unit normal bundle. It is the image of a *Fuchsian group* (that is, a discrete subgroup of  $\text{SL}_2(\mathbb{R})$ ) by a Lie group embedding of  $\text{SL}_2(\mathbb{R})$  in  $\text{PSU}_h$ . The extended  $\mathbb{C}$ -Fuchsian subgroups are then finite extensions of  $\mathbb{C}$ -Fuchsian subgroups by finite groups of complex reflections fixing the projective line or positive point or chain in the definition above. In particular, up to commensurability, the notions of extended  $\mathbb{C}$ -Fuchsian subgroups and of  $\mathbb{C}$ -Fuchsian subgroups coincide. The  $\mathbb{C}$ -Fuchsian lattices have been studied under a different viewpoint than our geometric one, as fundamental groups of arithmetic curves on ball quotient surfaces or Shimura curves in Shimura surfaces, by many authors, see for instance [16], [12], [13], [20] and their references.

An element of  $\Gamma_K$  is  *$K$ -irreducible* if it does not preserve a point or a line defined over  $K$  in  $\mathbb{P}_2(\mathbb{C})$ . An element of  $\mathbb{P}_2(\mathbb{C})$  is *rational* if it lies in  $\mathbb{P}_2(K)$ . Note that the polar line of a positive rational point of  $\mathbb{P}_2(\mathbb{C})$  is defined over  $K$ . The group  $\text{PSU}_h(K)$ , image of  $\text{SU}_h(K) = \text{SU}_h \cap \text{SL}_3(K)$  in  $\text{PSU}_h$ , preserves  $\mathbb{P}_2(K)$ , but in general its projective action on  $\mathbb{P}_2(K)$  is transitive neither on the positive, nor on the null, nor on the negative points of  $\mathbb{P}_2(K)$ .

The Galois group  $\text{Gal}(\mathbb{C} | K)$  acts on the complex projective plane  $\mathbb{P}_2(\mathbb{C})$  by  $\sigma[z_0 : z_1 : z_2] = [\sigma z_0 : \sigma z_1 : \sigma z_2]$ , and fixes  $\mathbb{P}_2(K)$  pointwise. Note that it does not preserve the positive, null, or negative cone of  $h$  in  $\mathbb{P}_2(\mathbb{C})$ . A positive point  $z \in \mathbb{P}_2(\mathbb{C})$  is *Hermitian cubic* over  $K$  if it is cubic over  $K$  (that is, if its orbit under  $\text{Gal}(\mathbb{C} | K)$  has exactly three points), and if its other Galois conjugates  $z', z''$  over  $K$  are null elements in the polar line of  $z$ .

The following result, analogous to [19, Prop. 9.6.1] in the Bianchi group case, strengthens one direction of [20, Lem. 1.2].



PROPOSITION 3.2. *A nonelementary extended  $\mathbb{C}$ -Fuchsian subgroup  $\Gamma$  of  $\Gamma_K$  fixes a unique rational point in  $\mathbb{P}_2(\mathbb{C})$ . This point is positive and it is the polar point of the unique complex geodesic preserved by  $\Gamma$ .*

PROOF. If  $\alpha \in \text{PSU}_h$  is loxodromic, let  $\alpha_-, \alpha_+ \in \partial_\infty \mathbb{H}_\mathbb{C}^2$  be its repelling and attracting fixed points, and let  $\alpha_0$  be its positive fixed point. Since the two projective lines tangent to the hypersphere  $\mathcal{HS}$  at  $\alpha_-$  and  $\alpha_+$  are invariant under  $\alpha$ , their unique intersection point is fixed by  $\alpha$ , therefore is equal to  $\alpha_0$ . In particular,  $\alpha_0$  is polar to the complex projective line through  $\alpha_-, \alpha_+$  (see also [21] for a more analytic proof).

Let  $L$  be the complex projective line preserved by  $\Gamma$ , which meets  $\mathbb{H}_\mathbb{C}^2$ . As  $\Gamma$  is nonelementary, there are loxodromic elements  $\alpha, \beta \in \Gamma$  such that their sets of fixed points in  $\partial_\infty \mathbb{H}_\mathbb{C}^2 \cap L$  are disjoint, see for instance [11, 8.2.E]. Since  $L$  passes through  $\alpha_-, \alpha_+$  as well as through  $\beta_-, \beta_+$ , and by the uniqueness of the polar point to  $L$ , we hence have  $\alpha_0 = \beta_0$ .

As  $\alpha$  and  $\beta$  have infinite order, one of them cannot be  $K$ -irreducible. Otherwise, if both were  $K$ -irreducible, then by [23, Prop. 20], the point  $\alpha_0 = \beta_0$  would be Hermitian cubic and its orbit under  $\text{Gal}(\mathbb{C}|K)$  would be  $\{\alpha_-, \alpha_+, \alpha_0\} = \{\beta_-, \beta_+, \beta_0\}$ , a contradiction. Assume then for instance that  $\alpha$  preserves a line or a point defined over  $K$ . As any projective subspace preserved by  $\alpha$  is a combination of  $\alpha_-, \alpha_+$  and  $\alpha_0$ , and as  $\alpha_-$  and  $\alpha_+$  are not defined over  $K$ , it follows that  $\alpha_0$  is rational.

Let  $\Gamma$  be a nonelementary extended  $\mathbb{C}$ -Fuchsian subgroup of  $\Gamma_K$ . By the previous proposition,  $\Gamma$  fixes a unique rational point  $P_\Gamma$  in  $\mathbb{P}_2(\mathbb{C})$ , which may be written  $P_\Gamma = [z_0 : z_1 : z_2]$  with  $z_0, z_1, z_2 \in \mathcal{O}_K$  relatively prime. Such a writing is unique up to the simultaneous multiplication of  $z_0, z_1, z_2$  by a unit in  $\mathcal{O}_K$ . Since the units in  $\mathcal{O}_K$  have norm 1, and since the trace and norm of  $K$  take integral values on the integers of  $K$ , the number

$$\Delta_\Gamma = h(z_0, z_1, z_2) = N(z_1) - \text{tr}(z_0 \bar{z}_2) \in \mathbb{Z}$$

is well defined, we call it the *discriminant* of  $\Gamma$  (by analogy with the case of  $\text{PSL}_2(\mathcal{O}_K)$ , see [18, Def. 4.1], and as it discriminates  $\Gamma$  up to finite error by Theorem 1.1). As  $P$  is positive, we have  $\Delta_\Gamma \in \mathbb{N} - \{0\}$ . The radius of the vertical projection of the polar chain of  $P_\Gamma$  is hence  $\frac{\sqrt{\Delta_\Gamma}}{|z_2|}$ . The discriminant of  $\Gamma$  depends only on the conjugacy class of  $\Gamma$  in  $\Gamma_K$ : for every  $\gamma \in \Gamma_K$ , since by uniqueness we have  $P_{\gamma\Gamma\gamma^{-1}} = \gamma P_\Gamma$ , we have

$$\Delta_{\gamma\Gamma\gamma^{-1}} = \Delta_\Gamma.$$

A chain  $C$  is ( $K$ -)arithmetic if its stabiliser in  $\Gamma_K$  has a dense orbit in  $C$ . The following result along with Proposition 3.2 justifies this terminology. This

result is well known, and it is the other direction of [20, Lem. 1.2], see also [12, Prop. 1.5, §III.1] and [16, §3]. We give a proof, which is a bit different, for the sake of completeness. In Section 4, we give an explicit construction based on quaternion algebras that implies this result.

**PROPOSITION 3.3.** *The stabiliser  $\text{Stab}_{\Gamma_K} P$  of any positive rational point  $P \in \mathbb{P}_2(K)$  is a maximal nonelementary extended  $\mathbb{C}$ -Fuchsian subgroup of  $\Gamma_K$ , whose invariant chain is arithmetic.*

**PROOF.** Let  $\underline{G}$  be the linear algebraic group defined over  $\mathbb{Q}$ , such that  $\underline{G}(\mathbb{Z}) = \text{PSU}_h(\mathcal{O}_K)$  and  $\underline{G}(\mathbb{R}) = \text{PSU}_h$ . We endow  $\mathbb{P}_2(\mathbb{C})$  with the  $\mathbb{Q}$ -structure  $\underline{X}$  whose  $\mathbb{Q}$ -points are  $\mathbb{P}_2(K)$  so that the action of  $\underline{G}$  on  $\underline{X}$  is defined over  $\mathbb{Q}$ .


As seen in Section 2, the set of real points of the stabilizer  $\text{Stab}_{\underline{G}} P$  is isomorphic to  $(\mathbb{S}^1 \times \text{SL}_2(\mathbb{R})) / \{\pm(1, \text{id})\}$  as a real Lie group. The group  $\text{Stab}_{\underline{G}} P$  is reductive and it has a (semisimple) Levi subgroup  $\underline{H}$  defined over  $\mathbb{Q}$ , such that  $\underline{H}(\mathbb{R})$  is isomorphic to  $\text{SL}_2(\mathbb{R})$ . By a theorem of Borel-Harish-Chandra [2, Theo. 7.8], the group  $\underline{H}(\mathbb{Z})$  is an arithmetic lattice in  $\underline{H}(\mathbb{R})$ , which (preserves the projective line polar to  $P$  and) is contained in  $\text{Stab}_{\Gamma_K} P$ . As  $\underline{H}(\mathbb{Z})$  is a lattice in  $\underline{H}(\mathbb{R})$ , the group  $\text{Stab}_{\Gamma_K} P$  is nonelementary and has a dense orbit in the chain  $P^\perp \cap \mathcal{HS}$ .

Recall that in the coordinates  $(w, -2 \text{Im } w_0)$  of  $\text{Heis}_3$ , the chains are ellipses whose images under the vertical projection are Euclidean circles (see also [10, §4.3]). The figure in the introduction is the vertical projection of part of the orbit under  $\Gamma_K$  of the chain  $[-5 : 0 : 1]^\perp \cap \mathcal{HS}$  when  $K = \mathbb{Q}[i]$ , so that  $\Gamma_K$  is the Gauss-Picard modular group, and we use the generating set of  $\Gamma_K$  given by [6]. The figure shows the square  $|\text{Re } z|, |\text{Im } z| \leq 1.5$  in  $\mathbb{C}$  with projections of chains whose diameter is at least 1.

In the figures on the following page (contrarily to the introduction where  $K = \mathbb{Q}(i)$ , so that the apparent symmetries are different),  $K = \mathbb{Q}[\omega]$ , where  $\omega$  is a primitive third root of unity, so that  $\Gamma_K$  is the Eisenstein-Picard modular group, and we use the generating set of  $\Gamma_K$  given by [8]. These two pictures illustrate the difference between  $D$  odd and  $D$  even in the coming proof of Theorem 1.1 (note in particular the difference around the origin).

The figure on the left shows part of the orbit of  $[-1 : 0 : 1]^\perp \cap \mathcal{HS}$  and the figure on the right shows part of the orbit of  $[-2 : 0 : 1]^\perp \cap \mathcal{HS}$ . They both show the square  $|\text{Re } z|, |\text{Im } z| \leq 1$  in  $\mathbb{C}$  with projections of chains whose diameter is at least 0.5 in the figure on the left and at least 0.75 in the figure on the right.

The first part of Theorem 1.1 in the introduction concerns the classification up to conjugacy in  $\Gamma_K$  of the maximal nonelementary extended  $\mathbb{C}$ -Fuchsian subgroups of  $\Gamma_K$ . Consider the set  $\mathcal{F}_{\mathbb{C}}$  of maximal nonelementary  $\mathbb{C}$ -Fuchsian



subgroups of  $\Gamma_K$ , on which the group  $\Gamma_K$  acts by conjugation. We will prove that the discriminant map  $\Gamma \mapsto \Delta_\Gamma$  on  $\mathcal{F}_\mathbb{C}$  induces a finite-to-one map from  $\Gamma_K \backslash \mathcal{F}_\mathbb{C}$  onto  $\mathbb{N} - \{0\}$ . Since every maximal nonelementary  $\mathbb{C}$ -Fuchsian subgroup  $\Gamma$  of  $\Gamma_K$  is contained in a unique maximal nonelementary extended  $\mathbb{C}$ -Fuchsian subgroup  $\widehat{\Gamma}$  of  $\Gamma_K$ , and since two maximal nonelementary  $\mathbb{C}$ -Fuchsian subgroups  $\Gamma, \Gamma'$  of  $\Gamma_K$  are conjugate if  $\widehat{\Gamma}, \widehat{\Gamma}'$  are conjugate, this implies the first part of Theorem 1.1.

The second part of Theorem 1.1 concerns the classification up to commensurability and conjugacy in  $\text{PSU}_h$ . Given a group  $G$  and a subgroup  $H$  of  $G$ , recall that two subgroups  $\Gamma, \Gamma'$  of  $H$  are *commensurable* if  $\Gamma \cap \Gamma'$  has finite index in  $\Gamma$  and in  $\Gamma'$ , and are *commensurable up to conjugacy in  $G$*  (or *commensurable in the wide sense*) if there exists  $g \in G$  such that  $\Gamma'$  and  $g\Gamma g^{-1}$  are commensurable. Two groups  $A$  and  $B$  are *abstractly commensurable* if they contain finite index subgroups  $A'$  and  $B'$  respectively that are isomorphic.

For any positive integer  $D$ , let

$$\Gamma_{K,D} = \text{Stab}_{\Gamma_K}[-D : 0 : 1].$$

The group  $\Gamma_{K,D}$  is, by Proposition 3.3, a maximal nonelementary extended  $\mathbb{C}$ -Fuchsian subgroup, which preserves the projective line  $[-D : 0 : 1]^\perp$ . Its discriminant is  $2D$ . We will prove that every element of  $\mathcal{F}_\mathbb{C}$  with discriminant  $D$  is commensurable up to conjugacy in  $\text{PSU}_h$  with  $\Gamma_{K,2D}$ .

PROOF OF THEOREM 1.1. (1) Let  $D \in \mathbb{N} - \{0\}$  and let

$$\mathcal{F}_\mathbb{C}(D) = \{\Gamma \in \mathcal{F}_\mathbb{C} : \Delta_\Gamma = D\}.$$

Let

$$P_D = \begin{cases} \left[-\frac{D}{2} : 0 : 1\right] & \text{if } D \text{ is even} \\ [0 : 1 : 0] & \text{if } D = 1 \\ \left[-\frac{D-1}{2} : 1 : 1\right] & \text{if } D > 1 \text{ is odd.} \end{cases}$$

By Proposition 3.3, the stabiliser in  $\Gamma_K$  of the positive rational point  $P_D$  is a maximal nonelementary extended  $\mathbb{C}$ -Fuchsian subgroup of  $\Gamma_K$ , with discriminant  $D$ . Hence  $\mathcal{F}_{\mathbb{C}}(D)$  is nonempty.

Let  $\underline{G}$  be the connected semisimple linear algebraic group defined over  $\mathbb{Q}$  such that  $\underline{G}(\mathbb{Z}) = \mathrm{SU}_h(\mathcal{O}_K)$  and  $\underline{G}(\mathbb{R}) = \mathrm{SU}_h$ . Let  $\pi : \underline{G} \rightarrow \mathrm{GL}(\underline{V})$  be the rational representation such that  $\underline{V}(\mathbb{Z}) = (\mathcal{O}_K)^3$ ,  $\underline{V}(\mathbb{R}) = \mathbb{C}^3$  and  $\pi|_{\underline{G}(\mathbb{R})}$  is the linear action of  $\mathrm{SU}_h$  on  $\mathbb{C}^3$ . Let  $\underline{X}_D$  be the closed algebraic submanifold of  $\underline{V}$  with equation  $h = D$ . In particular,  $\underline{X}_D$  is defined over  $\mathbb{Q}$ , and  $\underline{X}_D(\mathbb{R})$  is homogeneous under  $\underline{G}(\mathbb{R}) = \mathrm{SU}_h$ , by Witt's theorem. The map

$$\underline{X}_D(\mathbb{Z}) = \underline{X}_D \cap \underline{V}(\mathbb{Z}) = \{(z_0, z_1, z_2) \in (\mathcal{O}_K)^3 : h(z_0, z_1, z_2) = D\} \rightarrow \mathcal{F}_{\mathbb{C}},$$

which to  $(z_0, z_1, z_2)$  associates the stabiliser of  $[z_0 : z_1 : z_2]$  in  $\Gamma_K$  (which is the image of  $\underline{G}(\mathbb{Z})$  by the canonical map  $\underline{G}(\mathbb{R}) = \mathrm{SU}_h \rightarrow \mathrm{PSU}_h$ ), is well defined by Proposition 3.3 and  $\underline{G}(\mathbb{Z})$ -equivariant, and its image contains  $\mathcal{F}_{\mathbb{C}}(D)$ . Hence the finiteness of  $\Gamma_K \backslash \mathcal{F}_{\mathbb{C}}(D)$  follows from the finiteness of the number of orbits of  $\underline{G}(\mathbb{Z})$  on  $\underline{X}_D(\mathbb{Z})$ , see [2, Thm. 6.9].

(2) Let  $\Gamma \in \mathcal{F}_{\mathbb{C}}$ , and let  $D \in \mathbb{N} - \{0\}$  be its discriminant. By Propositions 3.2 and 3.3, and by maximality, there exists a unique positive rational point  $P = [z_0 : z_1 : z_2]$  with  $z_0, z_1, z_2$  relatively prime in  $\mathcal{O}_K$  such that  $\Gamma = \mathrm{Stab}_{\Gamma_K} P$  and  $D = h(z_0, z_1, z_2)$ .

*CLAIM.* *There exists  $\gamma \in \mathrm{PSU}_h(K)$  such that  $\gamma P = [-2D : 0 : 1]$ .*

Assuming this claim for the moment, we conclude the proof of the second part of Theorem 1.1: The groups  $\gamma \Gamma \gamma^{-1}$  and  $\Gamma_{K,2D}$  are commensurable, since

$$\gamma (\mathrm{Stab}_{\Gamma_K} P) \gamma^{-1} \cap \Gamma_{K,2D} = \mathrm{Stab}_{\gamma \Gamma_K \gamma^{-1} \cap \Gamma_K} \gamma P = \gamma (\mathrm{Stab}_{\Gamma_K \cap \gamma^{-1} \Gamma_K \gamma} P) \gamma^{-1}$$

and since  $\mathrm{PSU}_h(K)$  is the commensurator of  $\Gamma_K = \mathrm{PSU}_h(\mathcal{O}_K)$  in  $\mathrm{PSU}_h$  (see [1, Theo. 2]).

The following result, useful for the proof of the above claim, also gives a natural condition for when two such groups  $\Gamma_{K,D}$  for  $D \in \mathbb{N} - \{0\}$  are commensurable up to conjugacy in  $\mathrm{PSU}_h$ . A necessary and sufficient condition when  $D$  is even will be given in Corollary 4.2.

**LEMMA 3.4.** *If  $D, D' \in \mathbb{N} - \{0\}$  satisfy  $D' \in DN(\mathcal{O}_K)$ , then  $\Gamma_{K,D}$  and  $\Gamma_{K,D'}$  are commensurable up to conjugacy in  $\mathrm{PSU}_h(K)$ .*

PROOF. Let  $D \in \mathbb{N} - \{0\}$  and  $N \in N(\mathcal{O}_K) - \{0\}$ . As seen above, we only have to prove that there exists  $\gamma \in \text{PSU}_h(K)$  such that  $\gamma[-D : 0 : 1] = [-DN : 0 : 1]$ .

Assume first that  $D_K \equiv 0 \pmod{4}$ , so that  $\mathcal{O}_K = \mathbb{Z} + \frac{\sqrt{D_K}}{2}\mathbb{Z}$ . Since  $N \in N(\mathcal{O}_K)$ , there exists  $x, y \in \mathbb{Z}$  such that  $N = x^2 - \frac{D_K}{4}y^2$ . It is easy to check using Equation (1) and the fact that  $K = \mathbb{Q} + i\sqrt{|D_K|}\mathbb{Q}$  that the matrix

$$\gamma = \begin{pmatrix} x & 0 & -\frac{i}{2}\sqrt{|D_K|}Dy \\ 0 & 1 & 0 \\ -\frac{i}{2}\sqrt{|D_K|}\frac{y}{DN} & 0 & \frac{x}{N} \end{pmatrix}$$

belongs to  $\text{SU}_h(K)$ . Let  $\gamma$  be its image in  $\text{PSU}_h(K)$ . It is easy to check that as wanted  $\gamma[-D : 0 : 1] = [-DN : 0 : 1]$ .

If  $D_K \equiv 1 \pmod{4}$ , so that  $\mathcal{O}_K = \mathbb{Z} + \frac{1+\sqrt{D_K}}{2}\mathbb{Z}$ , the same argument works when  $\gamma$  in the above proof is replaced by the matrix

$$\begin{pmatrix} x + \frac{y}{2} & 0 & -\frac{i}{2}\sqrt{|D_K|}Dy \\ 0 & 1 & 0 \\ -\frac{i}{2}\sqrt{|D_K|}\frac{y}{DN} & 0 & \frac{x+\frac{y}{2}}{N} \end{pmatrix}$$

and the equation  $N = x^2 + xy + \frac{1-D_K}{4}y^2$  with  $x, y \in \mathbb{Z}$ .

PROOF OF THE CLAIM. As the lattice  $\Gamma_K$  does not preserve the complex geodesic with equation  $z_2 = 0$ , we may assume that  $z_2$  is nonzero, up to replacing  $P$  by an element in its orbit under  $\Gamma_K$ , which does not change the discriminant  $D$  of  $\Gamma$ . Let  $\gamma_1$  be the Heisenberg translation by the element

$$\left[ w_0 = \frac{|z_1|^2}{2|z_2|^2} - i \operatorname{Im} \frac{z_0}{z_2} : w = -\frac{z_1}{z_2} : 1 \right] \in \text{Heis}_3,$$

which belongs to  $\text{PSU}_h(K)$ . An easy computation shows that

$$\gamma_1[z_0 : z_1 : z_2] = [-D : 0 : 2N(z_2)].$$

Let  $\gamma_2$  be the image in  $\text{PSU}_h(K)$  of the diagonal element

$$\begin{pmatrix} 2N(z_2) & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \frac{1}{2N(z_2)} \end{pmatrix}$$

in  $\text{SU}_h(K)$ . Then  $\gamma_2\gamma_1$  maps  $P$  to  $[-2DN(z_2) : 0 : 1]$ . By the previous lemma, there exists  $\gamma_3 \in \text{PSU}_h(K)$  such that  $\gamma_3[-2DN(z_2) : 0 : 1] = [-2D : 0 : 1]$ . Hence the claim follows with  $\gamma = \gamma_3\gamma_2\gamma_1$ .

#### 4. Quaternion algebras

We refer to [28] and [19] for generalities on quaternion algebras. Let  $a, b \in \mathbb{Z}$  with  $a > 0$  and  $b < 0$ . The quaternion algebra  $A = \left(\frac{a,b}{\mathbb{Q}}\right)$  is the 4-dimensional central simple algebra over  $\mathbb{Q}$  with standard generators  $i, j, k$  satisfying the relations  $i^2 = a, j^2 = b$  and  $ij = -ji = k$ . The (reduced) norm of an element of  $A$  is

$$n(x_0 + x_1i + x_2j + x_3k) = x_0^2 - ax_1^2 - bx_2^2 + abx_3^2.$$

The group of elements in  $A(\mathbb{Z}) = \mathbb{Z} + i\mathbb{Z} + j\mathbb{Z} + k\mathbb{Z}$  with norm 1 is denoted by  $A(\mathbb{Z})^1$ . Let  $\sqrt{b}$  be the square root of  $b$  with positive imaginary part.

LEMMA 4.1. *The map  $\sigma = \sigma_{a,b} : A \rightarrow \mathcal{M}_3(\mathbb{C})$  defined by*

$$(x_0 + x_1i + x_2j + x_3k) \mapsto \begin{pmatrix} x_0 + x_1\sqrt{a} & 0 & (x_2 + x_3\sqrt{a})\sqrt{b} \\ 0 & 1 & 0 \\ (x_2 - x_3\sqrt{a})\sqrt{b} & 0 & x_0 - x_1\sqrt{a} \end{pmatrix}$$

*is a morphism of  $\mathbb{Q}$ -algebras and  $\sigma(A(\mathbb{Z})^1)$  is a discrete subgroup of the stabiliser of  $[0 : 1 : 0]$  in  $\mathrm{SU}_h$ .*

PROOF. It is well-known (and easy to check), see for instance [14], [19], that the map  $\sigma' : A \rightarrow \mathcal{M}_2(\mathbb{R})$  defined by

$$(x_0 + x_1i + x_2j + x_3k) \mapsto \begin{pmatrix} x_0 + x_1\sqrt{a} & (x_2 + x_3\sqrt{a})\sqrt{|b|} \\ -(x_2 - x_3\sqrt{a})\sqrt{|b|} & x_0 - x_1\sqrt{a} \end{pmatrix}$$

is a morphism of  $\mathbb{Q}$ -algebras and that the image of  $A(\mathbb{Z})^1$  is a discrete subgroup of  $\mathrm{SL}_2(\mathbb{R})$ . The map

$$\iota : \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mapsto \begin{pmatrix} a & 0 & ib \\ 0 & 1 & 0 \\ -ic & 0 & d \end{pmatrix}$$

is a morphism of  $\mathbb{Q}$ -algebras, sending  $\mathrm{SL}_2(\mathbb{R})$  into the stabiliser of  $[0 : 1 : 0]$  in  $\mathrm{SU}_h$  (see Equation (1)). The claim follows by noting that  $\sigma = \iota \circ \sigma'$ .

PROOF OF THEOREM 1.3. By Theorem 1.1, we only have to prove that the maximal  $\mathbb{C}$ -Fuchsian subgroup  $F_D$  of  $\Gamma_K$  stabilising  $[-2D : 0 : 1]$  (which has finite index in the extended  $\mathbb{C}$ -Fuchsian subgroup  $\Gamma_{K,2D}$ ) arises from the quaternion algebra  $\left(\frac{D, D_K}{\mathbb{Q}}\right)$ . It is easy to check that the element

$$\gamma_0 = -\frac{1}{\sqrt{2}} \begin{pmatrix} \sqrt{D} & \sqrt{2D} & \sqrt{D} \\ 1 & 0 & -1 \\ \frac{1}{2\sqrt{D}} & -\frac{1}{\sqrt{2D}} & \frac{1}{2\sqrt{D}} \end{pmatrix}$$

belongs to  $SU_h$  and maps  $[0 : 1 : 0]$  to  $[-2D : 0 : 1]$ . Hence, using Equation (1), a matrix  $M \in SU_h(\mathcal{O}_K)$  has its image (by the canonical projection  $SU_h \rightarrow PSU_h$ ) in  $F_D$  if and only if there exists  $a, d \in \mathbb{R}$  and  $b, c \in i\mathbb{R}$  with  $ad - bc = 1$  such that  $M = \gamma_0 \begin{pmatrix} a & 0 & b \\ 0 & 1 & 0 \\ c & 0 & d \end{pmatrix} \gamma_0^{-1}$ . A straightforward computation gives

$$M = \begin{pmatrix} \frac{1}{4}(a+b+c+d+2) & \frac{\sqrt{D}}{2}(a-b+c-d) & \frac{D}{2}(a+b+c+d-2) \\ \frac{1}{4\sqrt{D}}(a+b-c-d) & \frac{1}{2}(a-b-c+d) & \frac{\sqrt{D}}{2}(a+b-c-d) \\ \frac{1}{8D}(a+b+c+d-2) & \frac{1}{4\sqrt{D}}(a-b+c-d) & \frac{1}{4}(a+b+c+d+2) \end{pmatrix}.$$

This matrix has entries in  $\mathcal{O}_K$  if and only if

$$\begin{cases} a+b+c+d-2 \in 8D\mathcal{O}_K, \\ a+b-c-d \in 4\sqrt{D}\mathcal{O}_K, \\ a-b+c-d \in 4\sqrt{D}\mathcal{O}_K, \\ a-b-c+d \in 2\mathcal{O}_K. \end{cases}$$

Let  $u = a + d$ ,  $v = \frac{1}{2\sqrt{D}}(a - d)$ ,  $s' = b + c$  and  $t' = \frac{1}{2\sqrt{D}}(b - c)$ . Then  $M$  has entries in  $\mathcal{O}_K$  if and only if

$$\begin{cases} u + s' - 2 \in 8D\mathcal{O}_K, \\ v + t' \in 2\mathcal{O}_K, \\ v - t' \in 2\mathcal{O}_K, \\ u - s' \in 2\mathcal{O}_K. \end{cases} \quad (2)$$

Let  $D'_K = \frac{D_K}{4}$  if  $D_K \equiv 0 \pmod{4}$  and  $D'_K = D_K$  otherwise (so that  $K = \mathbb{Q}(\sqrt{D'_K})$ , see Section 3). Recall that  $\mathcal{O}_K \cap \mathbb{R} = \mathbb{Z}$  and  $\mathcal{O}_K \cap i\mathbb{R} = \mathbb{Z}\sqrt{D'_K}$ . The equations (2) imply in particular that  $u, v, s', t' \in \mathcal{O}_K$ . Note that  $a, d \in \mathbb{R}$  is equivalent to  $u, v \in \mathbb{R}$ , and  $c, b \in i\mathbb{R}$  is equivalent to  $s', t' \in i\mathbb{R}$ . Hence  $u, v \in \mathbb{Z}$  and there exists  $s, t \in \mathbb{Z}$  such that  $s' = s\sqrt{D'_K}$ ,  $t' = t\sqrt{D'_K}$ . Therefore

$$\gamma_0^{-1} F_D \gamma_0 = \left\{ \begin{pmatrix} \frac{u}{2} + v\sqrt{D} & 0 & (\frac{s}{2} + t\sqrt{D})\sqrt{D'_K} \\ 0 & 1 & 0 \\ (\frac{s}{2} - t\sqrt{D})\sqrt{D'_K} & 0 & \frac{u}{2} - v\sqrt{D} \end{pmatrix} : \begin{array}{l} u, v, s, t \in \mathbb{Z} \\ v + t\sqrt{D'_K} \in 2\mathcal{O}_K \\ v - t\sqrt{D'_K} \in 2\mathcal{O}_K \\ u - s\sqrt{D'_K} \in 2\mathcal{O}_K \\ u + s\sqrt{D'_K} - 2 \in 8D\mathcal{O}_K \end{array} \right\}.$$

The group  $\gamma_0^{-1}F_D\gamma_0$  is contained in  $\sigma_{D,D'_K}(A(\mathbb{Z})^1)$ , since the parameters  $u$  and  $s$  have to be even as a consequence of the defining equations of  $\gamma_0^{-1}F_D\gamma_0$ . Furthermore,  $\gamma_0^{-1}F_D\gamma_0$  contains  $\sigma_{D,D'_K}(\mathcal{O}^1)$ , where  $\mathcal{O}$  is the order of  $A$  defined by

$$\mathcal{O} = \{x_0 + ix_1 + jx_2 + kx_3 \in A(\mathbb{Z}) : x_1, x_2, x_3 \equiv 0 \pmod{4D}\}.$$

Indeed, if  $x_0 + ix_1 + jx_2 + kx_3 \in \mathcal{O}^1$ , then with  $u = 2x_0, s = 2x_2, v = x_1, t = x_3$ , we have, since  $x_0 \equiv 1 \pmod{4D}$  by the condition  $n(x_0 + ix_1 + jx_2 + kx_3) = 1$ ,

$$\begin{cases} v \pm t\sqrt{D'_K} \in 2\mathbb{Z} + 2\sqrt{D'_K}\mathbb{Z} \subset 2\mathcal{O}_K \\ u - s\sqrt{D'_K} \in 2\mathbb{Z} + 2\sqrt{D'_K}\mathbb{Z} \subset 2\mathcal{O}_K \\ u - 2 + s\sqrt{D'_K} = 2(x_0 - 1) + 2x_2\sqrt{D'_K} \in 8D\mathbb{Z} + 8D\sqrt{D'_K}\mathbb{Z} \subset 8D\mathcal{O}_K. \end{cases}$$

Since  $\sigma_{D,D'_K}(\mathcal{O}^1)$  has finite index in  $\sigma_{D,D'_K}(A(\mathbb{Z})^1)$  (see for instance [28], Coro. 1.5 in Chapt. IV), the groups  $\gamma_0^{-1}F_D\gamma_0$  and  $\sigma_{D,D'_K}(A(\mathbb{Z})^1)$  are commensurable.

Since  $\left(\frac{D,D'_K}{\mathbb{Q}}\right) = \left(\frac{D,D_K}{\mathbb{Q}}\right)$  as  $D'_K$  and  $D_K$  differ by a square factor, the result follows.

Observe that, by Theorem 1.3, a maximal nonelementary  $\mathbb{C}$ -Fuchsian subgroup of  $\Gamma_K$  of discriminant  $D$  is cocompact (in its copy of  $\mathrm{SL}_2(\mathbb{R})$ ) if and only if  $\left(\frac{D,D_K}{\mathbb{Q}}\right)$  is a division algebra (see for instance [14, Thm. 5.4.1]).

**COROLLARY 4.2.** *Let  $D, D' \in \mathbb{N} - \{0\}$ . The subgroups  $\Gamma_{K,2D}$  and  $\Gamma_{K,2D'}$  are commensurable up to conjugacy in  $\mathrm{PSL}_h$  if and only if the quaternion algebras  $\left(\frac{D,D_K}{\mathbb{Q}}\right)$  and  $\left(\frac{D',D_K}{\mathbb{Q}}\right)$  are isomorphic.*

**PROOF.** We have seen in the previous proof that  $\Gamma_{K,2D}$  arises from the quaternion algebra  $\left(\frac{D,D_K}{\mathbb{Q}}\right)$ . The result hence follows from the fact that two arithmetic Fuchsian groups are commensurable up to conjugacy in  $\mathrm{SL}_2(\mathbb{R})$  if and only if their associated quaternion algebras are isomorphic (see [27]).

The following corollaries follow from the arguments in [17], pages 309 and 310. Corollary 1.2 of the introduction follows from Corollary 4.4 below. We refer for instance to [19] for the unexplained terminology below.

**PROPOSITION 4.3.** *Let  $A$  be an indefinite quaternion algebra over  $\mathbb{Q}$ . There exists an arithmetic  $\mathbb{C}$ -Fuchsian subgroup of  $\Gamma_K$  whose associated quaternion algebra is  $A$  if and only if the primes at which  $A$  is ramified are either ramified or inert in  $K$ .*



**COROLLARY 4.4** (Chinburg-Stover). *Every Picard modular group  $\Gamma_K$  contains infinitely many wide commensurability classes in  $\text{PSU}_h$  of maximal non-elementary  $\mathbb{C}$ -Fuchsian subgroups.*

**COROLLARY 4.5.** *Any arithmetic Fuchsian group whose associated quaternion algebra is defined over  $\mathbb{Q}$  has a finite index subgroup isomorphic to a  $\mathbb{C}$ -Fuchsian subgroup of some Picard modular group  $\Gamma_K$ .*

**COROLLARY 4.6.** *For all quadratic irrational number fields  $K$  and  $K'$ , there are infinitely many abstract commensurability classes of arithmetic Fuchsian subgroups with representatives in both Picard modular groups  $\Gamma_K$  and  $\Gamma_{K'}$ .*

#### REFERENCES

1. Borel, A., *Density and maximality of arithmetic subgroups*, J. Reine Angew. Math. 224 (1966), 78–89.
2. Borel, A., and Harish-Chandra, *Arithmetic subgroups of algebraic groups*, Ann. of Math. (2) 75 (1962), 485–535.
3. Cartan, E., *Sur le groupe de la géométrie hypersphérique*, Comment. Math. Helv. 4 (1932), no. 1, 158–171.
4. Chinburg, T., and Stover, M., *Fuchsian subgroups of lattices acting on hermitian symmetric spaces*, preprint arxiv:1105.1154v3.
5. Chinburg, T., and Stover, M., *Geodesic curves on Shimura surfaces*, preprint arxiv:1506.03299.
6. Falbel, E., Franciscs, G., and Parker, J. R., *The geometry of the Gauss-Picard modular group*, Math. Ann. 349 (2011), no. 2, 459–508.
7. Falbel, E., and Parker, J. R., *The moduli space of the modular group in complex hyperbolic geometry*, Invent. Math. 152 (2003), no. 1, 57–88.
8. Falbel, E., and Parker, J. R., *The geometry of the Eisenstein-Picard modular group*, Duke Math. J. 131 (2006), no. 2, 249–289.
9. Fine, B., *Algebraic theory of the Bianchi groups*, Monographs and Textbooks in Pure and Applied Mathematics, vol. 129, Marcel Dekker, Inc., New York, 1989.
10. Goldman, W. M., *Complex hyperbolic geometry*, Oxford Mathematical Monographs, The Clarendon Press, Oxford University Press, Oxford, 1999.
11. Gromov, M., *Hyperbolic groups*, Math. Sci. Res. Inst. Publ., vol. 8, pp. 75–263, Springer-Verlag, New York, 1987.
12. Holzapfel, R.-P., *Arithmetic curves on ball quotient surfaces*, Ann. Global Anal. Geom. 1 (1983), no. 2, 21–90.
13. Holzapfel, R.-P., *Ball and surface arithmetics*, Aspects of Mathematics, E29, Friedr. Vieweg & Sohn, Braunschweig, 1998.
14. Katok, S., *Fuchsian groups*, Chicago Lectures in Mathematics, University of Chicago Press, Chicago, IL, 1992.
15. Knapp, A. W., *Lie groups beyond an introduction*, Progress in Mathematics, vol. 140, Birkhäuser Boston, Inc., Boston, MA, 1996.
16. Kudla, S. S., *Intersection numbers for quotients of the complex 2-ball and Hilbert modular forms*, Invent. Math. 47 (1978), no. 2, 189–208.
17. Maclachlan, C., *Fuchsian subgroups of the groups  $\text{PSL}_2(O_d)$* , in *Low-dimensional topology and Kleinian groups* (Coventry/Durham, 1984), London Math. Soc. Lecture Note Ser., vol. 112, Cambridge Univ. Press, Cambridge, 1986, pp. 305–311.

18. Maclachlan, C., and Reid, A. W., *Parametrizing Fuchsian subgroups of the Bianchi groups*, *Canad. J. Math.* 43 (1991), no. 1, 158–181.
19. Maclachlan, C., and Reid, A. W., *The arithmetic of hyperbolic 3-manifolds*, *Graduate Texts in Mathematics*, vol. 219, Springer-Verlag, New York, 2003.
20. Möller, M., and Toledo, D., *Bounded negativity of self-intersection numbers of Shimura curves in Shimura surfaces*, *Algebra Number Theory* 9 (2015), no. 4, 897–912.
21. Parker, J. R., *Complex hyperbolic Kleinian groups*, Cambridge Univ. Press, to appear.
22. Parkkonen, J., and Paulin, F., *Prescribing the behaviour of geodesics in negative curvature*, *Geom. Topol.* 14 (2010), no. 1, 277–392.
23. Parkkonen, J., and Paulin, F., *Counting and equidistribution in Heisenberg groups*, *Math. Ann.* 367 (2017), no. 1-2, 81–119.
24. Samuel, P., *Théorie algébrique des nombres*, Hermann, Paris, 1967.
25. Serre, J.-P., *Lie algebras and Lie groups*, second ed., *Lecture Notes in Mathematics*, vol. 1500, Springer-Verlag, Berlin, 1992.
26. Stover, M., *Volumes of Picard modular surfaces*, *Proc. Amer. Math. Soc.* 139 (2011), no. 9, 3045–3056.
27. Takeuchi, K., *A characterization of arithmetic Fuchsian groups*, *J. Math. Soc. Japan* 27 (1975), no. 4, 600–612.
28. Vignéras, M.-F., *Arithmétique des algèbres de quaternions*, *Lecture Notes in Mathematics*, vol. 800, Springer-Verlag, Berlin, 1980.
29. Zimmer, R. J., *Ergodic theory and semisimple groups*, *Monographs in Mathematics*, vol. 81, Birkhäuser Verlag, Basel, 1984.

DEPARTMENT OF MATHEMATICS AND STATISTICS  
P.O. BOX 35  
40014 UNIVERSITY OF JYVÄSKYLÄ  
FINLAND  
*E-mail*: jouni.t.parkkonen@jyu.fi

LABORATOIRE DE MATHÉMATIQUE D'ORSAY  
UMR 8628 UNIV. PARIS-SUD ET CNRS  
UNIVERSITÉ PARIS-SACLAY  
91405 ORSAY CEDEX,  
FRANCE  
*E-mail*: frederic.paulin@math.u-psud.fr